

ANTIOXIDANT ACTIVITY OF *FICUS CARICA* AND *PRUNUS AMYGDALUS* VAR. *DULCIS* – A COMPARATIVE STUDY

*Naazia Zafar

Centre for Pharmaceutical Sciences, Jawaharlal Nehru Technological University, Hyderabad, Telangana, India.

*Corresponding Author: Naazia Zafar M.Pharm

Centre for Pharmaceutical Sciences, Jawaharlal Nehru Technological University, Hyderabad, Telangana, India.

Article Received on 27/09/2017

Article Revised on 18/10/2017

Article Accepted on 08/11/2017

ABSTRACT

Antioxidants are the essential molecules or substances which possess the potential to prevent or protect the human body from destruction caused by free radical impelled oxidative stress. Epidemiological studies indicate that increase intake of fruits and vegetables have the ability to hinder the damaging behavior of free radicals in the human body. In this study, we analyzed the *in - vitro* antioxidative activity of aqueous extracts of commonly used dry fruits like *Ficus carica* L. and *Prunus amygdalus* var. *dulcis* Mill. The extracts were analyzed for antioxidant activity by Hydroxyl radical scavenging method and were compared to standard and commonly used antioxidant Ascorbic acid (AA). The concentration required for 50% inhibition (IC₅₀ - Inhibitory Concentration) was estimated. All the extracts showed potential free radical scavenging activity which increased with increasing concentration. The analysis was made with the use of UV-Vis Spectrophotometer (Model Shimadzu UV-1800) at a wavelength of 532 nm. The present study revealed that Hydroxyl radical scavenging activity of *Ficus carica* (IC₅₀ value of 2064 µg/ml) was more than that of *Prunus amygdalus* var. *dulcis* (IC₅₀ value of 7274 µg/ml). Multiple flavonoids and related polyphenols contribute crucially to the total antioxidant activity as well as therapeutic uses of many fruits and vegetables. Natural dietary antioxidants with health benefits like Figs and Almonds are preferred because synthetic antioxidants are considered to be carcinogenic.

KEYWORDS: Antioxidant activity, free radicals, *Ficus carica*, *Prunus amygdalus* var. *dulcis*, Ascorbic acid.**INTRODUCTION****Aim**

Human body has inherent mechanism to reduce the free radical induced injury by endogenous antioxidants. Sometimes these protective mechanisms were found to be not sufficient when compared to the insult produced to the body. Hence, the search for exogenous antioxidants is continued.

Review of Literature

An antioxidant is a molecule that has the ability to prevent the oxidation of other molecules. Oxidation is a chemical reaction that transfers electrons or hydrogen from a substance to an oxidizing agent. These reactions can produce free radicals. In turn, these radicals can start chain reactions. When the chain reaction occurs in a cell, it can cause damage or death to the cell.

Antioxidants stop or end these chain reactions by removing free radical intermediates and inhibit other oxidation reactions.^[1] They do this by being oxidized themselves, so they are often known as reducing agents. Plants and animals have the capability to maintain complex systems of multiple types of antioxidants such

as glutathione, vitamin C, vitamin A, and vitamin E as well as enzymes such as catalase, superoxide dismutase and various peroxidases.^[2] Insufficient levels of antioxidants or inhibition of the antioxidative enzymes causes oxidative stress and may damage or kill cells. Oxidative stress is damage to cell wall, cell structure, cell function and genetic material by overly reactive oxygen-containing molecules or reactive oxygen species (ROS)^[3] like superoxide anion, singlet oxygen, hydroxyl radical and hydrogen peroxide.

The oxidative damage initiated by ROS is propagated by lipid peroxidation which may cause damage to DNA. Oxidative stress seems to play a significant role in many human diseases, including cancers, AIDS, atherosclerosis, stroke, coronary heart disease, diabetes, arthritis, neurodegenerative diseases, liver cirrhosis, aging, altitude sickness etc.^[4]

Table 1: Classification of Endogenous Antioxidants.^[5]

Category	Examples
Low Molecular Weight Proteins	Lipid soluble: Tocopherols, Carotenoids, Quinones, Bilirubin Water soluble: Ascorbic acid, Uric acid.
High Molecular Weight Proteins	Albumin, Ceruloplasmin, Transferrin, Haptoglobin.
Enzymes	Superoxide dismutase, Catalase, Glutathione peroxidase.

Mechanism of action of Antioxidants^[6]

- Electron donation: Primary antioxidants are compounds which are able to donate hydrogen atom rapidly to a lipid radical, forming a new radical, more stable than the initial one. Biological organs contain many polyunsaturated fatty acids (PUFA) such as linoleic, linolenic and arachidonic acids, mainly in the form of esters of phospholipids, triglycerides or with cholesterol. These PUFA can undergo lipid peroxidation which can be interrupted by antioxidants by the donation of electrons.
- Metal Chelation: Secondary antioxidants can retard the rate of radical initiation reaction by means of initiators elimination. This can be accomplished by deactivation of high energy species (e.g. singlet oxygen), absorption of UV light, scavenging of oxygen and thus reducing its concentration, chelation of metal catalyzing free radical reaction or by inhibition of peroxidases such as NADPH oxidase, xanthine oxidase, dopamine-hydroxylase or lipoxygenase.
- Co – antioxidants: Ascorbic acid alone has little effect in preventing lard oil from oxidation. The combination of ascorbic acid with tocopherol gave rise to a strong synergistic antioxidative effect. The role of ascorbic acid was to preserve tocopherol from consumption. This behaviour of ascorbic acid is termed as co - antioxidant effect.
- Gene Expression: Antioxidant possesses the ability to donate electrons and thereby act as reducing agents, to chelate metal ions and thereby remove potential radical initiators and to facilitate

antioxidant activity by other compounds (co – antioxidants). Antioxidants can also effect directly or indirectly the expression of genes in tissues. A number of genes are regulated by changes in the cellular redox status.

Screening of Antioxidants^[7,8,9]

Antioxidants are screened by the following methods

- Superoxide radical/Riboflavin Photoreduction method.
- Hydroxyl radical/Deoxyribose method.
- Peroxide radical / Induction of Fe⁺² / ascorbate system method.
- DPPH (2, 2-diphenyl-1-picrylhydrazyl) scavenging / Quenching activity method.
- Trolox equivalent antioxidant capacity (TEAC) method / ABTS (2, 2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) radical cation decolorization assay.
- Nitric oxide scavenging activity.
- Potassium ferricyanide reduction method.
- Total radical-trapping antioxidant parameter (TRAP) method.
- Hydroxyl radical averting capacity (HORAC) method.
- Oxygen radical absorbance capacity (ORAC) method.

Table 2: Natural Sources of Antioxidants^[10,11,12]

Antioxidant Compound	Sources
Dry fruits	Figs, almonds, raisins, prunes, walnut, cashew nut, pistachio, dates, apricot, pecans.
Vitamin C	Fresh fruits and vegetables.
Vitamin E	Cereals, nuts, vegetable oils.
Polyphenols (Flavonoids)	Turmeric, tea, coffee, soy, olive oil, chocolate, cinnamon, oregano, apples, grapes, berries.
Carotenoids	Carrot, beet root, capsicum, tomato, leak, red cabbage, kale
Green leafy vegetables	Spinach, cabbage, mint, parsley, celery, broccoli, peppers, chillies, Brussels sprouts.
Spices	Cloves, pepper, coriander, cumin, dill, fennel, caraway.

OBJECTIVE OF THE RESEARCH^[13,14,15,16]

Figs and Almonds both have polyphenols, which is responsible for their antioxidant activity. Dry Figs have 193 (mg/g) whereas Almonds have 114 (mg/g) of total phenolic content which is expressed in terms of Gallic acid equivalent (GAE). The objective of the research was to determine the antioxidant activity or free radical scavenging activity of Figs (*Ficus carica* L., Family:

Moraceae) and Almonds (*Prunus amygdalus* var. *dulcis* Mill., Family: Rosaceae) using *in - vitro* method of Hydroxyl radical assay and comparing their activity with that of a well-established antioxidant, Ascorbic acid.

MATERIALS AND METHODS

Dry figs and Almonds were purchased from local market of Hyderabad, India. Ascorbic acid, Ferric chloride and

Ethylenediaminetetraacetic acid (EDTA) were purchased from Prime Laboratories, Hyderabad. 2 – Deoxy – D – ribose was a gift sample from SRL Pvt. Ltd, Hyderabad. Disodium phosphate and Sodium dihydrogen phosphate were purchased from S. D. Fine Chemicals, Mumbai, India. Chemicals of analytical grade and deionized water were used throughout the study.

Preparation of Extracts^[17]

500 gm. dry fruits of Figs were washed with water to remove the dirt and later, dried and cut into small pieces. The dried pieces of Figs were grinded and macerated with 500 ml of water for 1 hour. Later, the mixture was transferred to a round bottomed flask and was boiled for ½ hour attached to a condenser by which the material was never allowed to get dried. After boiling, the mixture was filtered. The filtrate obtained was heated to reduce its volume to half of its original volume and later kept for evaporation by air. In the same way, the extraction of dry Almonds was carried out. The crude extracts were re-dissolved in distilled water, as and when necessary for the assessment of antioxidant activity.

Preparation of Reagents

- 37.54 mg of 2 – Deoxy – D – ribose was weighed, transferred to a volumetric flask and the volume was made up to 10 ml with distilled water to give 28mM solution. From this 0.1 ml was used in the reaction.
- 37.22 mg of EDTA was weighed, transferred to a volumetric flask and the volume was made up to 100 ml with distilled water to give 1mM solution. From this 0.1 ml was used in the reaction.
- 16.22 mg of Ferric Chloride was weighed, transferred to a volumetric flask and the volume was made up to 100 ml with distilled water to give 1mM solution. From this 0.1 ml was used in the reaction.
- 17.66 mg of Ascorbic acid was weighed, transferred to a volumetric flask and the volume was made up to 100 ml with distilled water to give 1mM solution. From this 0.1 ml was used in the reaction.
- Phosphate buffer solution (PBS), P^H 7.4:
Solution A: 276 mg of Disodium phosphate (Na₂HPO₄ 2H₂O) was weighed into a beaker and the volume was made up to 100 ml with distilled water.

Solution B: 568 mg of Sodium dihydrogen phosphate (NaH₂PO₄ 2H₂O) was weighed into a beaker and the volume was made up to 100 ml with distilled water.

From the above, 12 ml of solution A and 88 ml of solution B were mixed and the P^H was adjusted to 7.4.

Hydroxyl Radical Assay (Deoxyribose Method)^[18,19]

Hydroxyl radical scavenging activity of the extract was found by studying the competition between deoxyribose and the test compounds i.e. extract for the hydroxyl radical generated from Fe⁺³ / ascorbate / EDTA / H₂O₂

system. Hydroxyl radical acts against deoxyribose which results in the formation of thiobarbituric acid reacting substance (TBARS).

Procedure

Preparation of Test Sample

For the experiment, the extracts of dry Figs and Almonds were used in the concentrations of 2000µg, 4000µg, 6000µg and 8000µg each. To 0.4 ml of phosphate buffer solution (PBS), 0.1 ml of extract solution was added. To the above mixture, 0.1 ml of EDTA, 0.1 ml of FeCl₃, 0.1 ml of deoxyribose, 0.1 ml of vitamin C and 0.1 ml of H₂O₂ were added and the reaction mixture was incubated at 37° C. From the above 1 ml reaction mixture, 0.4 ml was withdrawn and treated with 0.2 ml dodecyl sulphate (8.1 %), 1.5 ml thiobarbituric acid (0.8 %) and 1.5 ml acetic acid (20 %, P^H 3.5). The total volume was then made up to 4 ml by adding distilled water and kept in oil bath maintained at 95° C for 1 hour. After cooling, the absorbance was measured at 532 nm (Spectrophotometer Model Shimadzu UV – 1800).

The same above experiment was repeated with Ascorbic acid as test sample, in the same concentrations of 2000µg, 4000µg, 6000µg and 8000µg as that of the extracts.

Preparation of Control Sample

Control sample is prepared by the same procedure as that of test sample without the addition of extract / ascorbic acid.

The percentage inhibition was calculated by comparing the optical density of each test sample with that of optical density of control sample by using the below formula

$$\text{Percentage inhibition} = \frac{\text{Control sample O.D} - \text{Test sample O.D}}{\text{Control sample O.D}} \times 100$$

The mean and standard error of the mean of five samples of each concentration of Figs, Almonds and Ascorbic acid were calculated and presented in Table 3. The data was plotted by taking concentration on X - axis and percentage inhibition on Y - axis as shown in Fig. 1, 2 and 3. From the graph, the concentration required for the 50 % inhibition of hydroxyl radicals was obtained and the results are shown in Table 4. The concentrations of Figs, Almonds and Ascorbic acid required for 50 % inhibition were also represented by bar diagram as shown in Fig. 4.

RESULTS

Table 3: Hydroxyl radical scavenging activity of *Ficus carica* and *Prunus amygdalus* var. *dulcis* and Ascorbic acid.

S. No.	Extract / Drug	Concentration ($\mu\text{g} / \text{ml}$)	Optical Density	Percentage Inhibition
1.	<i>Ficus carica</i>	2000	0.388	48.40% \pm 0.26
		4000	0.023	96.94% \pm 0.82
		6000	-	-
		8000	-	-
2.	<i>Prunus amygdalus</i> var. <i>dulcis</i>	2000	0.648	13.82% \pm 0.41
		4000	0.545	27.52% \pm 0.35
		6000	0.442	41.22% \pm 0.19
		8000	0.339	54.92% \pm 0.64
3.	Ascorbic acid	2000	0.599	20.34% \pm 0.59
		4000	0.445	40.82% \pm 0.49
		6000	0.300	60.10% \pm 0.71
		8000	0.141	81.25% \pm 0.95





Figure 1: Percentage Inhibition of *Ficus carica*.Figure 2: Percentage Inhibition of *Prunus amygdalus* var. *dulcis*.

Figure 3: Percentage Inhibition of Ascorbic acid.

Table 4: Concentration ($\mu\text{g} / \text{ml}$) at which 50 % of hydroxyl radicals were inhibited.

S. No.	Extract / Drug	50% Inhibition was obtained at Conc ($\mu\text{g}/\text{ml}$)
1.	<i>Ficus carica</i>	2064
2.	<i>Prunus amygdalus</i> var. <i>dulcis</i>	7274
3.	Ascorbic acid	4937

Figure 4: Concentration ($\mu\text{g} / \text{ml}$) required for 50 % Inhibition.

DISCUSSION

Degradation of deoxyribose mediated by hydroxyl radicals generated by Fe^{+3} / ascorbate / EDTA / H_2O_2 system was found to be initiated by both *F. carica* and *P. amygdalus* var. *dulcis* extracts. The extracts of *F. carica* (at quantities 2000 μg and 4000 μg) and *P. amygdalus* var. *dulcis* (at quantities 2000 μg , 4000 μg , 6000 μg and 8000 μg) scavenged the hydroxyl radicals in a dose dependent manner. Ascorbic acid at concentrations of 2000 μg , 4000 μg , 6000 μg and 8000 μg was also found to produce dose dependent inhibition of hydroxyl radicals. The quantity of *F. carica* extract required to produce 50 % inhibition by hydroxyl radicals was 2064 μg whereas that of *P. amygdalus* var. *dulcis* extract was 7274 μg . Similar effect was produced by Ascorbic acid at concentration of 4937 μg .

CONCLUSION

The present study indicated strong antioxidant activity of the aqueous extract of *Ficus carica* dry fruits which was 2.3 – 2.4 times more than that of Ascorbic acid, a well-established and therapeutically used antioxidant. But the antioxidant activity of aqueous extract of *Prunus amygdalus* var. *dulcis* dry fruits was found to be 0.6 – 0.7 times less than that of Ascorbic acid, which was very less when compared to *Ficus carica*. Strong antioxidant activity of Figs when compared to Almonds is due to its higher total phenolic content. The presence of such polyphenol compounds which are responsible for their antioxidant activity in *in vitro* studies, might lead to their use in Ayurveda for beneficial effects, in treatment of certain disorders.

ACKNOWLEDGEMENT

The author is thankful to JNTU, Hyderabad for providing the facilities to carry out this study.

REFERENCES

- Borg DC. Oxygen Free Radicals and Tissue Injury. In: Tarr M and Samson F (eds.). Oxygen Free Radicals in Tissue Damage. Massachusetts, Birkhäuser Boston, 1993; 12-55.
- Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn Rev*, 2010; 4(8): 118–126.
- Apel K, Hirt H. Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu Rev Plant Biol*, 2004; 55: 373-399.
- Pham-Huy LA, He H, Pham-Huy C. Free radicals, antioxidants in disease and health. *Int J Biomed Sci*, 2008; 4(2): 89–96.
- Li J, Liu X, Shen L, Zeng W, Qiu G. Natural plant polyphenols for alleviating oxidative damage in man, current status and future perspectives. *Trop J Pharm Res*, 2016; 15(5): 1089-1098.
- Brewer MS. Natural antioxidants: Sources, compounds, mechanisms of action and potential applications. *Comp Rev in Food Sci F Saf*, 2011; 10(4): 221–247.
- Fukai T, Ushio-Fukai M. Superoxide dismutases: Role in redox signaling, vascular function and diseases. *Antioxid Redox Signal*, 2011; 15(6): 1583–1606.
- Kramarenko GG, Hummel SG, Martin SM, Buettner GR. Ascorbate reacts with singlet oxygen to produce Hydrogen peroxide. *Photochem Photobiol*, 2006; 82(6): 1634–1637.
- Alam MN, Bristi NJ, Rafiquzzaman M. Review on *in vivo* and *in vitro* methods evaluation of antioxidant activity. *Saudi Pharm J*, 2013; 21(2): 143-152.
- Vijaya Kumar Reddy C, Sreeramulu D, Raghunath M. Antioxidant activity of fresh and dry fruits commonly consumed in India. *Food Res Int*, 2010; 43(1): 285-288.
- Kusum Devi V, Rehman F. Nutraceutical Antioxidants - An overview. *Ind J Pharma Edu*, 2002; 36(1): 3-8.
- Krinsky NI. The antioxidant and the biological properties of the carotenoids. *Ann NY Acad Sci*, 1998; 854: 443-447.
- Dr Nadkarni KM. *Indian Materia Medica*. 3rd ed., Vol. I. Mumbai; Popular Prakashan, 1954; 545- 547, 1036, 1037.
- Trease GE, Evan WC. *Textbook of Pharmacognosy*. 16th ed., Oxford; Alden Press, 2009; 217, 348.
- Kamiloglu S, Capanoglu E. Polyphenol content in Figs (*Ficus carica* L.): Effect of sun-drying. *Int J Food Prop*, 2015; 18(3): 521-535.
- Abe LT, Lajolo FM, Genovese MI. Comparison of phenol content and antioxidant capacity of nuts. *Food Sci Tech*, 2010; 30(1): 254-259.
- Biren N Shah, Seth AK. *Textbook of Pharmacognosy and Phytochemistry*. 1st ed., Gurgaon; Elsevier India, 2010; 410.
- Elizabeth K, Rao MNA. Oxygen radical scavenging activity of curcumin. *Int J Pharm*, 1990; 58: 237-240.
- Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Ana Biochem*, 1979; 95(2): 351-358.