

DISTRIBUTION OF ANTIBIOTIC RESISTANT BACTERIA FLORA IN READY TO-EAT FOOD SAMPLES SOLD IN CALABAR METROPOLIS

Ashang U.J., Antai S.P. and Tiku D.R.

Department of Microbiology, University of Calabar, Calabar.

***Corresponding Author: Ashang U.J.**

Department of Microbiology, University of Calabar, Calabar.

Article Received on 19/10/2017

Article Revised on 09/11/2017

Article Accepted on 30/11/2017

ABSTRACT

The distribution of antibiotic resistant bacteria flora in ready-to-eat food samples sold in Calabar metropolis was investigated. Thirty (30) samples of different ready-to-eat food samples (cooked indomie, sausage rolls, egg-roll, bean cake, fried meat and fish) were purchased at the point of sales in various eateries located in University of Calabar campus, Watt Market and Marian market and were placed into sterile specimen bottles with screw caps, labeled and transported immediately to the laboratory in an ice-packed container for further analysis. The study was completed within a duration of six months. All the procedures were carried out using standard microbiological techniques. The results of the study revealed the presence of *Pseudomonas spp*, *Escherichia spp*, *Bacillus spp*, *Staphylococcus aureus*, *Enterobacter spp*, *Klebsiella spp*, *Proteus spp* and *Shigella spp*. *Escherichia coli* had the highest percentage of occurrence (17.83%) compared to other bacterial isolate in the analyzed ready-to-eat food samples that had; *Klebsiella spp* (15.50%), *Salmonella spp* (14.72%), *Enterobacter spp* (6.98%), *Proteus spp* (10.85%), *Shigella spp* (9.30%), *Bacillus spp* (4.65%), *Pseudomonas spp* (6.89%) and *Staphylococcus aureus* (6.20%). Bacteria isolates from the analyzed ready-to-eat food samples showed a higher percentage resistance to augmentin (100%) and amoxicillin (100%) compared to other antibiotics tested against Cotrimoxazole (55.56%) nitrofurantoin (66.67%), gentamycin (66.67%) ofloxacin (44.44%), tetracycline (55.56%), ciprofloxacin (77.78%) and nalidixic acid (22.22%). Among the bacteria isolates from the ready-to-eat food samples, *Staphylococcus aureus* had the highest percentage resistant to the antibiotics tested against (Ceftriazone, amoxicillin, cotrimoxazole, nitrofurantoin, augmentin and tetracycline) while *Klebsiella spp* and *Shigella spp* showed sensitivity to all the antibiotics tested against. However, the study has revealed the distribution and occurrence of multiple antibiotic resistance among bacteria isolates in ready-to-eat food samples sold in Calabar Metropolis. Thus, there is need for good sanitary practices, and intensive surveillance of isolates throughout food pre- and post production stages so as to reduce or eliminate cases of food borne infections, as well as to also detect emerging antimicrobial resistance bacterial phenotypes.

KEYWORDS: Antibiotics, Resistant, Antimicrobial, Bacterial, Food-borne, Infections.**INTRODUCTION**

Anti-microbial resistant bacteria have been recovered from both healthy humans (Irona *et al*, 2012) and a wide variety of food which include vegetables (Mora *et al*, 2005), confectionary (Okeke *et al*, 2005), meat and meat products and poultry (Schoeder *et al*, 2004). Food and water contaminated by faecal material from healthy human may also be an important source of antibiotic resistant organisms that later cause human infections (Schoeder *et al*, 2004).

Contamination of ready to eat food may occur during and after processing, and this is of primary concern because such organisms may be pathogenic thereby leading to outbreak of food-borne illness (Russel and Path, 2001). Moreover, non-pathogenic organism that can contaminate man's food chain from time to time may

serve as reservoir of genes for antimicrobial resistance in organisms (Goldstein *et al*, 2001).

A myriad of substances such as drugs radionuclide and solvents are used in hospital, for medical treatment, diagnostics, disinfection and research (Diab *et al* 2008), and after application many non-metabolized drugs mainly antibiotics are excreted by the patients and from there it enters waste water and then other environment. Antibiotics are whole range of chemical substances that kill or inhibit the growth of bacteria (Brook *et al*, 2008), most are naturally produced by living organisms while others are produced synthetically (Brook *et al*, 2008). They are selectively toxic (affecting pathogenic microorganism more adversely than the host), and this may be as a result of the function of specific receptors required for drugs attachment or it may depend on the inhibition of biochemical events essential to the pathogen

but not to the host (King *et al.*, 2000). The most selective agents of antibiotics are those affecting structures (e.g. cell wall) or functions (e.g. folic acid synthesis) present only in prokaryotic cells (Walsh and Howe, 2002), while the less selective antibiotics are those affecting protein or nucleic acid synthesis which are essential functions for both prokaryotic (bacteria) and eukaryotic cells (the host) (Walsh and Howe, 2002).

Antibiotics when in these waste water may be present at levels that can not only alter the ecology of the environment but also give rise to antibiotic resistance (Diab *et al.*, 2008). Acquired resistance to antibiotics may arise by cellular mutation or by acquisition of genetic elements in the form of plasmids or transposons (Diab *et al.*, 2008). The occurrence of strongly selective environments such as water and food promotes, not only the growth of resistant bacteria, but also leads to an increase in the frequency of resistant bacterial genes and genetic elements such as plasmids (Khachatourians *et al.*, 2003). Waste water when not effectively treated may contain pathogenic drug-resistant bacteria which constitute the most dangerous single risk factor for dissemination of pathogenic and drug resistant bacterial species in the environment (Cabrera *et al.* 2004). These resistant bacteria species may be transmitted to humans and farm animals hereby causing infection that cannot be treated by conventional antibiotics (Chitris, 2004). Therefore, this forms the main rationale why this study is focused on investigating the distribution of antibiotic resistant bacteria flora in ready-to-eat-foods in Calabar metropolis.

MATERIALS AND METHODS

The study area

The study was carried out in Calabar, which is the capital of Cross River State, Nigeria. It is bounded at north by Odukpani and Akpabuyo Local Government Area, at east by the Republic of Cameroun, at the west by Akwa-Ibom State, and the south by the Atlantic Ocean. Calabar with an approximate population of about one million two hundred thousand (1,200,000) inhabitants (2006 census), is situated seventy-seven (77) kilometers up the Calabar river and over an area of about 604km².

Calabar has an average high temperature of 29⁰c and an average low temperature of 25⁰c. The precipitation of Calabar is about 51mm on the average during the dry season and 445mm on average during the wet season (weather base, 2011). Calabar is located approximately between longitude 80⁰19¹E and 80⁰21¹E and latitude 40⁰55¹N and 40⁰58¹N. Calabar inhabited people are from three (3) ethnic groups which are the Quas, Efuts and the Efiks.

Sample collections

Ready-to-eat-food samples

30 samples of different ready-to-eat-foods comprising of moi-moi (beans pudding), indomie, meat and fish, sausage rolls, egg-roll, doughnut and bean cake were

purchased at the point of sales in various eateries located in University of Calabar Campus, Watt market and Marian market and were placed into sterile specimen bottles with tight screw caps, labeled and transported immediately to the laboratory in an ice-packed container for further analysis.

Media

The media used in this study were Nutrient agar, Muller Hinton agar, Motility Indole Ornithine (MIO) (Hardy diagnostics, USA); MacConkey agar, Simmon Citrate Medium (Acumedia, USA). These media were prepared according to the manufacturers instruction.

Chemical and reagents

Chemical used in this study were of analytical grade. They include absolute alcohol, acetone, methanol (Sigma, USA) neutral red, methyl red indicators, phenol red indicator (Titan Biotech, India). Reagents used were oxidase strips, indole kovacs and were products of Hardy diagnostics, USA.

Sample preparation

Ready-to-eat-food samples

10 grams of the ready-to-eat-food samples were aseptically weighed into 90ml of sterile distilled water in a 100ml conical flask. The sample was vortexed to homogenized and allowed to stand for 10minutes. From this initial dilution, 10-fold serial dilutions were carried out in clean sterile test tubes containing 9ml of sterile distilled water.

Plaiting procedures

0.1ml of desire dilutions 10⁻³ – 10⁻⁵ was spread plated in duplicate into nutrient agar and MacConkey agar plates supplemented with 50µg/ml of nystatin Plates were incubated at 37⁰c for 24hours bacterial counts were then recorded.

Purification of Isolates

Following enumeration of total heterotrophic bacteria, representatives of observed colonies were selected and sub-cultured repeatedly into nutrient agar for purification. Purified isolates were stocked in nutrient agar slants for further studies.

Characterization and identification

Purified bacterial isolates were characterized by gram staining and biochemical tests using the scheme in Bergey's manual of determinative bacteriology (Holt *et al.*, 1994; Cheesbrough, 2000).

Antibiotic susceptibility testing

The antibiotic susceptibility test was determined by the disc diffusion method as described by Bauer *et al.* (1996). Ten different commercially prepared antibiotic discs (Abtek Biological Ltd, Uk) were used and the concentration of each is as follows; amoxicillin (25µg), cotrimoxazde (25µg), nalidixic acid (30µg), nitrofurntoin (30µg), gentamycin (10µg), ofloxacin (30µg),

augmentin (30µg), tetracycline (30µg), doxycycline (30µg), trimethoprin/sulfamethazole (25µg).

After 18hours incubation of the isolates in Mueller Hinton agar at 37^oc, the size of the zone of inhibition was measured and interpreted by comparing with 0.5 Mcfarland standard antibiotic sensitivity chart to determine their resistance patterns.

RESULTS

Biochemical characterization and identification of isolate

bacteria isolates from the ready-to-eat food samples were identified as *Escherichia coli*, *pseudomonas spp*, *Enterobacter spp*, *Klebsiella spp*, *Salmonella spp*, *Bacillus spp*, *Staphylococcus aureus*, *Proteus spp* and *Shigella spp*,

Frequency and percentage occurrence of bacteria isolate from the ready-to-eat food samples

Table 1 presents the result of frequency of occurrence of bacteria isolate from ready-to-eat food and water samples respectively. It showed that *Escherichia coli* had the highest frequency of occurrence (23) compared to other isolate counterparts from ready-to-eat food samples that had; *Klebsiella spp*, (20), *Salmonella spp* (19), *Enterobacter spp* (18), *Proteus spp* (14), *Shigella spp* (12), *Pseudomonas spp* (9), and *Bacillus spp* (6).

Fig 1 presents the result of percentage occurrence of bacteria isolate from ready-to-eat food samples, it showed that *Escherichia coli* had the highest percentage occurrence (17.83%), compared to other isolates that had; *Klebsiella spp* (15.50%), *Salmonella spp* (14.72%), *Enterobacter spp* (13.95%), *Proteus spp* (10.85%), *Shigella spp* (9.30%), *Pseudomonas spp* (6.98%), *Staphylococcus aureus* (6.20%) and *Bacillus spp* (4.65%).

Antibiotic susceptibility profile of bacteria isolates from analyzed the ready-to- eat food sample

Table 2 presents the result of antibiotic susceptibility profile of bacteria isolate from the analyzed ready-to-eat food sample.

It showed that bacteria isolate from the ready-to-eat food samples. Its showed a higher percentage resistance to augmentin (100%) and amoxicillin (100%) compared to other isolates tested against; cotrimoxazole (55.56%), nitrofurantoin (66.67%), gentamycin (66.67%), ofloxacin (44.44%), tetracycline (55.56%), ciprofloxacin (77.78%) and ceftriazone (77.78%) while the bacteria isolate showed the least percentage resistance to nalidixic acid (22.22%).

Distribution and proportion of antibiotic resistance among bacterial isolates from samples

Table 3 presents the result of the distribution and proportion of antibiotic resistance among bacterial isolates from ready-to-eat food samples. It showed that

67.44% of the isolates showed resistance to ceftracazone, 83.72% showed resistance to amoxicillin, 50.39% showed resistance to cotrimoxazole, 55.81% showed resistance to nitrofurantoin, 68.22% showed resistance to gentamycin, 24.81% showed resistance to Nalidixic acid, 10.85% showed resistance to ofloxacin, 64.34% showed resistance to augmentin, 79.07% showed resistance to tetracycline, and 59.69% showed resistance to ciprofloxacin.

Fig 2 presents the result of antibiotics resistance pattern in *Escherichia coli* from ready-to-eat food samples. It showed that the isolate had a higher percentage of occurrence (100%) when tested against amoxycillin, while the isolate was all sensitive to ofloxacin compared other antibiotics tested against.

Fig 3 presents the result of antibiotic resistance pattern in *Klebsiella Spp* from ready-to-eat food samples. It showed that the isolate had a higher percentage of resistance with Ciprofloxacin (85%) and least resistance to ofloxacin and cotrioxazole (20% each).

Fig 4 presents the result of antibiotic resistance pattern in *Proteus Spp* from ready-to-eat food samples. It showed that the isolate showed a higher percentage resistance to ceftriazone (100%), amoxycillin (100%) and augmentin (100%), while the least resistance was observed with nalidixic acid (21.42%) and ofloxacin (21.42%) compared to other antibiotic.

Fig 5 presents the result of antibiotic resistance pattern in *Pseudomonas Spp* from ready-to-eat food samples. It showed that the isolate had a higher percentage resistance to ceftriazone (100%), amoxycillin (100%) and Tetracycline (100%), while the least percentage resistance was observed with ofloxacin (55.67%).

Fig 6 presents the result of antibiotic resistance pattern in *Salmonella Spp* from ready-to-eat food samples. It showed that the isolate had a higher percentage resistance to tetracycline (100%), and least percentage resistance were all sensitive to Ofloxacin compared to other antibiotics tested against.

Fig 7 presents the result of antibiotic resistance pattern in *Enterobacter Spp* from ready-to-eat food samples. It showed that the isolate had a higher percentage resistance when tested against Amoxycillin (100%), Gentormycin (100%) and Tetracycline (100%), and were all sensitive to Nalidixic acid, compared to other antibiotics tested against.

Fig 8 presents the result of antibiotic resistance pattern in *Shigella Spp* from ready-to-eat food samples. It showed that the isolate had a higher percentage resistance when tested against cotrimoxazole (83.33%) and Tetracycline (83.33%) and were all sensitive To ofloxacin as compared to other antibiotics tested against.

Fig 9 presents the result of antibiotic resistance pattern in *Bacillus Spp* from ready-to-eat food samples. It showed that the isolate had a higher percentage resistance to ceftriazone (100%), amoxicillin (100%) and least percentage resistance to ofloxacin (33.33%), as compared to other antibiotic tested against.

Table 1: Frequency of occurrence of bacteria isolate from ready-to-eat food samples.

Isolate	Frequency	% Occurrence
<i>Escherichia coli</i>	23	17.83
<i>Enterobacter spp</i>	18	13.95
<i>Staphylococcus aureus</i>	8	6.20
<i>Klebsiella spp</i>	20	15.50
<i>Salmonella spp</i>	19	14.73
<i>Pseudomonas spp</i>	9	6.89
<i>Proteus spp</i>	14	10.85
<i>Shigella spp</i>	12	9.30
<i>Bacillus spp</i>	6	4.65
Total	129	100




Fig. 1: Percentage occurrence of bacteria isolates from ready-to-eat-food samples.

Table 2: Antibiotic susceptibility profile of bacteria isolate from ready-to-eat food sample.

Antibiotics tested	Disc potency (µg/ml)	<i>Escherichia coli</i>	<i>Pseudomonas spp</i>	<i>Enterobacter spp</i>	<i>Klebsiella spp</i>	<i>Salmonella spp</i>	<i>Bacillus spp</i>	<i>Staphylococcus aureus</i>	<i>Proteus spp</i>	<i>Shigella</i>	% Resistance of organism
Amoxicillin	25	R	R	R	R	R	R	R	R	R	100
Cotrimoxazole	25	S	R	S	S	R	R	R	R	S	55.56
Nitrofurantoin	30	S	R	R	S	R	R	R	R	S	66.67
Gentamycin	10	R	R	R	S	R	R	R	S	S	66.67
Nalidixic acid	30	S	R	S	S	S	S	R	S	S	22.22
Ofloxacin	30	S	R	S	S	S	R	R	R	S	44.44
Augmentin	30	R	R	R	R	R	R	R	R	R	100
Tetracycline	30	S	R	S	S	R	R	R	R	S	55.56
Ciprofloxacin	10	S	S	R	R	R	R	R	R	R	77.78
Ceftriazone	30	S	R	R	R	R	R	R	R	S	77.78
% Resistance of single organisms		30	90	60	40	80	90	100	80	30	

Table 3: Distribution and proportion of antibiotic resistance among bacterial isolate from ready-to-eat food samples.

Isolates identified	No.	Total number and percentage resistance exhibited to an antibiotics									
		Ce (%)	AMX (%)	COT (%)	NIT (%)	GEN (%)	NAL (%)	OFL (%)	AUG (%)	TET (%)	CIP (%)
<i>Escherichia Coli</i>	23	10(43.48)	23(100%) fdaf 23(100)	7(30.4)	6(26.10)	21(91.30)	2(8.70)	0(0)	22(95.70)	8(34.50)	15(65.22)
<i>Klebsiella spp</i>	20	13(65)	12(60)	11(55)	4(20)	5(25)	3(21.40)	0(0)	4(20)	11(55)	17(85)
<i>Proteus spp</i>	14	14(100)	14(100)	10(71.43)	13(92.90)	10(71.40)	3(21.40)	3(21.42)	14(100)	13(92.9)	11(78.57)
<i>Pseudomonas spp</i>	9	9(100)	9(100)	9(100)	9(100)	8(88.90)	8(88.90)	5(55.67)	8(88.9)	9(100)	5(55.56)
<i>Salmonella spp</i>	19	12(63.16)	10(52.63)	6(31.58)	10(52.63)	10(52.63)	5(26.32)	0(0)	7(36.78)	19(100)	4(21.05)
<i>Enterobacter spp</i>	18	10(55.56)	18(100)	0(0)	9(50)	18(100)	0(0)	0(0)	8(44.44)	18(100)	7(38.89)
<i>Shigella spp</i>	12	5(41.67)	8(66.67)	10(83.33)	7(58.33)	5(41.67)	2(16.67)	0(0)	6(50)	10(83.3)	7(58.33)
<i>Bacillus spp</i>	6	6(100)	6(100)	4(66.67)	6(100)	4(66.67)	3(50)	2(33.33)	6(100)	6(100)	5(83.33)
<i>Staphylococcus aureus</i>	8	8(100)	8(100)	8(100)	8(100)	7(87.5)	6(75)	4(50)	8(100)	8(100)	6(75)
Total	129	87(64.44)	108(83.72)	65(50.39)	72(55.81)	88(68.22)	32(24.81%)	14(10.85%)	83(64.34%)	102(79.07)	77(59.69%)

AMX = Amoxicillin (25µg), COT = Cotrimoxazole (25µg), NIT = Nitrofurantoin (30µg), GEN = Gentamicin (10µg), NAL = Nalidixic acid (30µg), OFL = Ofloxacin (30µg), AUG = Augmentin (30µg), TET = Tetracycline (30µg), Ce = Ceftriazone (30µg), CIP = Ciprofloxacin (10µg)




Fig. 2: Antibiotic resistance pattern in *Escherichia coli* from ready-to-eat food samples.




Fig 5: Antibiotic resistance pattern in *Pseudomonas spp* from ready-to-eat food samples.




Fig 3: Antibiotic pattern in *Klebsiella sp* from ready-to-eat food samples.




Fig 6: Antibiotic resistance pattern in *Salmonella* from ready-to-eat food samples.




Fig 4: Antibiotic resistance pattern in *Proteus spp* from ready-to-eat food samples.




Fig 7: Antibiotic resistance pattern in *Enterobacter spp* from ready-to-eat food samples.




Fig. 8: Antibiotic resistance pattern in *Shigella spp* from ready-to-eat food samples.




Fig. 9: Antibiotic resistance pattern in *Bacillus spp* from ready-to-eat food samples.

DISCUSSION

In this study, bacteria isolate from ready-to-eat food and water samples were identified as *Escherichia coli*, *Pseudomonas Spp*, *Enterobacter spp*, *Klebsiella spp*, *Salmonella spp*, *Bacillus spp*, *Staphylococcus aureus*, *Proteus spp*, and *Shigella spp*. This observation was not surprising as it corroborates with that of Oluyeye *et al* 2009; Oladipo and Adejemobi, 2010; and Majolagbe *et al*, 2011, who all identified *Staphylococcus aureus*, *Bacillus*, *Marcescens*, *Streptococcus faccalis*, *Pseudomonas putida*, *Aeromonas hydrophila*, *Enterobacter aerogenes*, *Klebsiella spp*, and *Proteus spp* from ready-to-eat food samples in Ado-ekiti and Ogbomoso, Nigeria respectively.

Escherichia coli was the most common bacteria isolate from the collected ready-to-eat and samples with a percentage occurrence of 17.83% as compared to other isolate from the samples. This observation corroborates with that of Oluyeye *et al.*, (2009), who reported a higher occurrence of *Escherichia coli* in ready-to-eat food samples analyzed. *Escherichia coli* when found in water and food supplies, is an indicative of a recent faecal contamination and is a threat to public health (Moro *et*

al, 2000; Mora *et al*, 2005). Its presence is a major health concern especially in cases of verotoxin producing *E. coli* (VTEC) serogroup O157, a major cause of haemorrhagic colitis (Moro *et al.*, 2000). Faecal contamination of food and water cannot be prevented entirely, particularly in this setting where hygienic standard of food production and water treatment process is low and not monitored (Famurewa *et al.*, 2003). The frequency of isolation of *Salmonella* and *Shigella spp* was also reasonably high and supports individual studies and laboratory records that have however revealed that typhoid fever is endemic in Nigeria (Moro *et al.*, 2001). The percentage occurrence of *Klebsiella spp*, *Shigella spp* and *Enterobacter spp* were a bit high, their contamination could be attributed to the poor hygiene practices of the food handlers before and after food processes upon returning from toilet, lack of disinfection of table tops before and after daily use by customers as well as poor water treatment process (Fang *et al.*, 2013) as this postulation is supported by previous report on the isolation of *Salmonella spp*, *Klebsiella spp*, *Shigella spp*, *Staphylococcus aureus* and *Escherichia coli* from the hands of food vendors and food canteens in Nigeria (Famurewa *et al.*, 2003).

Antibiotics susceptibility profile of bacteria isolates from the ready-to-eat food samples showed that all the isolates had multiple resistance to the antibiotics they were tested against. This observation was in agreement with that of Gundogan *et al.*, (2006) who reported to have isolated *Klebsiella spp* and *Escherichia coli* with multiple antibiotic resistance from meat, chicken and meat ball samples. Also other studies by Lateef *et al.*, (2005); Majolagbe *et al.*, 2011, have reported to isolate *E. coli*, *Staphylococcus aureus*, *Salmonella spp* and *Streptococcus spp* from water, food and clinical samples with multiple antibiotic resistance.

The relatively high level of resistance of the isolates to antimicrobial agents, is a reflection of misuse or abuse of these agents in the environment (Abbar and Kaddar, 2001). Antibiotics prescriptions in hospitals are given without clear evidence of infection or adequate medical attention. Broad spectrum antibiotics sometimes given in place of narrow spectrum are drugs as a substitute for culture and sensitivity testing, with consequential risk being super infections and the selection of drug resistant mutants (Prescott *et al.*, 2005). In developing countries, drugs are available to the public and thus people may practice self-administration of antibiotics and further increase the prevalence of drug-resistant strains, also the long standing practice of using low doses of antibiotics for a long period of time for growth promotion and arbitrary use of antibiotics in animal husbandry is a strong contributor to the development of antibiotic resistant bacteria in the environment.

The ability of some of the isolates such as *Bacillus spp*, *Pseudomonas spp*, *Enterbacter spp*, *Salmonella spp*, *Proteus spp*, *E. coli* and *Staphylococcus aureus* to show

100% resistance to some of the antibiotics such as ceftriaxone, amoxicillin, augmentin, tetracycline and cotrimoxazole could possibly be as a result of selection pressure created by the use of antimicrobials in food producing animals as well as the indiscriminate use of antibiotics by human (Chui *et al* 2002; Threfall *et al.*, 2000). The co-existence of resistance genes with mobile elements such as plasmids, transposons and intergrons facilitates the rapid spread of antibiotic resistance genes among bacteria and this could possibly be the reason for the high percentage resistance (Sunde, 2005).

The total resistance (100%) shown by some of the isolates to ceftriaxone. (a cephalosporin and widely used as broad spectrum antibiotic) which acts by inhibiting cell wall synthesis in growing or dividing cells (Kathleen and Arthur, 2000) is likely due to the presence of β -Lactamase which acts by cleaving β -Lactam ring of cell wall, thus inhibiting antibiotics like ceftriaxone (Warren 2006) while the resistance to other antibiotics by the isolates could as well be due to the fact that antibiotic resistance microorganisms may be associated with reduced penetration of antibiotic into the cell, or from active processes such as changes in the transport of those compounds into or from the bacteria cells (Hermansson *et al.*, 2007).

Most of the bacteria isolates from the analyzed ready-to-eat samples showed significantly higher ($p < 0.05$) percentage resistance to the antibiotics tested. This observation was not surprising, as this could have possibly resulted from inappropriate or uncontrolled use of antibiotic in farming practices, indiscriminate use of antibiotics by humans, indiscriminate dumping of hospital wastes and antibiotic materials in the environment, indiscriminate use of manure as well as human excreta in the environment (Warburton *et al* 2002).

CONCLUSION

The study has revealed the distribution and occurrence of multiple antibiotic resistance among bacteria isolates in ready-to-eat food samples sold in Calabar Metropolis. The study thus emphasizes the need for intensive surveillance of isolates throughout the food production continuum as to prevent food borne infections, as well as detect emerging antimicrobial resistance bacterial phenotypes especially in this our developing world.

REFERENCES

1. Abbar, F., & Kaddar, H. Bacteriological studies on Iraq milk products. *Journal of Applied Bacteriology*, 2001; 71: 497–503.
2. Brook, L., Shaw, A. Sharp, D. & Hay, A. Towards a better understanding of patient's perspective of antibiotic resistance and MRSA. *Family Practice*, 2008; 25: 341-348.
3. Cabrera, M., Mangel, A. & Johnson, J. And distribution of urinary tract infection by a multiple drug resistant *Escherichia coli* clonal group. *Journal of Microbiology*, 2004; 1007-1013.
4. Chui, C., Wu, L., Chu, J., Chia, A., & Lin, Y. The emergence in Taiwan of Fluor quinolone resistance in *Salmonella enterica* serotype. *England Journal of Medicine*, 2002; 346: 416–423.
5. Chitris, L. Flow of resistance gene in the environment and from animals to man. *Journals of Microbial Chemotherapy*, 2004; 18: 189-189.
6. Cheesbrough, M. District laboratory practice in tropical countries (part 2). Cambridge University Press, London, 2000; 132-134.
7. Diab, A., Jeannette, M., Lang, K. & Lapara, T. Luating the effect of chlorotetracycline O, the proliferation of antibiotic resistance bacterial in stipulated river water ecosystem. *Applied and Environmental Microbiology*, 2008; 73(7): 5421-5425.
8. Famurewa, O., Oyagade, J., Femi-Ola, T. & Laleye, S. Assessment of Microbial quality of ready-to-eat food retailed in Ado – Ekiti. *BISEB Journal*, 2003; 3: 71–77.
9. Ghazi, M. & Rajaa, A. Studies on *Pseudomonas aeruginosa* in drinking water from water suppliers consumed in Basra. *Journal of Applied Microbiology*, 2014; 5: 17–26.
10. Gundogan, N., Devren, A. & Citak, S. Incidence, protease activity and antibiotic resistance of *Escherichia Coli* and *Klebsiella Spp* isolated from meat, chicken and meatball samples. *Journal of Medical Microbiology*, 2006; 57: 113–117.
11. Hermansson, M., Johnston, I. & Jaykus, L. Incident of Virulence factor and antibiotic resistance among microbes from food. *Applied Environmental Microbiology*, 2007; 67: 4385–4395.
12. Goldstein, C., Margie, D., Sanchez, S. & Hudson, C. Incidence of class 1 and 2 integrases in clinical and commensal bacteria from livestock, companion animals and exotic. *Antimicrobial Agents Chemotherapy*, 2001; 45: 723-726.
13. Khachatourians, Y., Kummerer, K. & Henniger, A. Antibiotic resistance by the emission from hospital and household into clinical microbial infection. *Applied Journal of Microbiological*, 2003; 32(3): 131-138.
14. Kathleen, S. & Arthur, D. Emergence and dissemination of antibiotic resistant *Escherichia coli* in the community. *Antimicrobial Agents Chemotherapy*, 2000; 43: 2736–2741.
15. King, D Malone, R. & Lilley, S. New classification and update on the quinone antibiotics. *American Family Physician*, 2000; 61: 2741-2748.
16. Lateef, A., Oloke, K. & Guegium, E. The prevalence of bacterial resistance in clinical food, water and some environmental samples in southwest Nigeria. *Environmental Monitoring and Assessment*, 2005; 100: 59–69.
17. Moro, D, Oluduro, A., Salu, O. & Famurewa, O. The prevalence of bacterial pathogens and intestinal

- worms among food vendors in Ajegunle, Lagos. *Journal of Biological Science.*, 2000; 1: 129–134.
18. Moro, D., Famurewa, O. & Agboola, A. *Salmonella* and Intestinal worms in food handlers in Ado-Ekiti, Nigeria. *Journal of Clinical Microbiology*, 2001; 27: 731–739.
 19. Majolagbe, O., Idowu, S., Adebayo, E., Ola, L., Adewoyin, A. & Oladipo, E. Prevalence and antibiotic resistance of bacteria isolated from ready-to-eat food (RTE) samples of highly patronized eateries in Ogbomoso – Oyo State, Nigeria. *Pelagia Research Library*, 2011; 1(3): 70–78.
 20. Mora, A., Blanco, J., Blanc, M. & Bernardes, M. Antimicrobial resistance of shiga toxin (verotoxin) producing *Escherichia coli* 0157:47 and non-0157 strains isolated from humans, cattle, sheep and food in Spain. *Research in Microbiology*, 2005; 3: 1-14.
 21. Oluyeye, A., Dada, A., Ojo, M. & Oluwadare, E. Antibiotic resistance profile of bacterial isolates from food sold on a university campus in south western Nigeria. *African Journal of Biotechnology*, 2009; 8(12): 5883–5887.
 22. Oladipo, I. & Adejumobi, O. Incidence of Antibiotic resistance in some bacterial pathogens from street vended food in Ogbomoso, Nigeria. *Parkhistian Journal of Nutrition*, 2010; 9(11): 1061–1068.
 23. Okeke, I. Lamikanra, A. & Steinruck, H. Characterization of *Escherichia coli* strains from cases of childhood diarrhea in provincial southwest Nigeria. *Journal of Clinical Microbiology*, 2000; 38: 7-12.
 24. Piddock, L. Clinically relevant chromosomally encoded multi-drug resistance efflux pumps in bacteria. *Clinical Microbiology Review*, 2006; 19: 382-402.
 25. Prescott, L., Harles, J. & Klein, D. *Microbiology*, 4th ed. McGraw Hill. New York, 2001; 678–697.
 26. Schoeder, C., White, D., & Meng, J. Retail meat and poultry as a reservoir of antimicrobial resistant *Escherichia Coli*. *Food Microbiology*, 2004; 21: 244–255.
 27. Sunde, M. Prevalence and characterization of class 1 and class 2 integrons in *Escherichia coli* isolated from meat and meat products of Norwegian origin. *Journal of Antimicrobial Chemotherapy*, 2005; 56: 1019–1027.
 28. Threefall, E., Ward, J., Frost, A. & Wilshaw, G. The emergence and spread of antibiotic resistance in foodborne bacteria. *International Journal of Food Microbiology*, 2000; 62: 1–5.
 29. Russel, A., & Path, R. Mechanism of bacterial susceptibility to biocide. *American Journal of Infection Control*, 2001; 29: 259-261.
 30. Warren, T. Veterinary use and antibiotic resistance. *Current Opinion in Microbiology*, 2006; 4: 493–499.
 31. Warburton, D., Girrafa, G. & Fram, C. Antimicrobial resistance of microorganisms isolated from food product. *Applied Environmental Microbiology*, 2002; 70: 3133–3139.
 32. Walsh, L & Howe, A. A study of the effect of the incidence of antibiotic resistance of antibiotic resistance in *E. coli*. *Biomedical Letters*, 2002; 55: 91-97.