

**HISTOPATHOLOGY STUDY TO PROVE THE PROTECTIVE EFFECT OF MAJORANA HORTENSIS AGAINST OXIDATIVE STRESS**Radha Palaniswamy<sup>1\*</sup> and Padma Raghunathan<sup>2</sup><sup>1</sup>Department of Biotechnology, Dr. NGP Arts and Science College, Kalapatti, Coimbatore - 641048.<sup>2</sup>Department of Biochemistry, Biotechnology and Bioinformatics, Avinashilingam University for Women, Coimbatore - 641043.**\*Corresponding Author: Dr. Radha Palaniswamy**

Department of Biotechnology, Dr. NGP Arts and Science College, Kalapatti, Coimbatore - 641048.

Article Received on 21/01/2018

Article Revised on 11/02/2018

Article Accepted on 05/03/2018

**ABSTRACT**

*Majorana hortensis* leaves have been proved to have potential antioxidant activity which was further proved by *in vitro* studies using precision cut liver slices. Histopathological examinations of the liver slices were used as to examine the hepatic architecture after the exposure of oxidant, namely H<sub>2</sub>O<sub>2</sub>. The microscopic lesions produced were usually fatty metamorphosis or centrilobular necroses which were studied. The group which consisted of oxidant and methanolic extract of the leaves showed a protective effect compared to the group which was exposed to the oxidant alone.

**KEYWORDS:** *Majorana hortensis*, oxidative stress, liver slices, hydrogen peroxide.**INTRODUCTION**

*Majorana hortensis* (*M.hortensis*) is a Mediterranean perennial herb of Lamiaceae family. Commonly it is called sweet majoram and used to add flavor in culinary purpose. It has been proved to have high potential as an antioxidant.<sup>[1]</sup> An alternate *in vitro* source such as precision cut goat liver slices were used in the study which was examined histopathologically. It helps to envisage the anatomy and gives the insight into the functioning of tissues.<sup>[2]</sup> The live slices were chosen as it plays a key role in the metabolism and biochemical transformations of oxidants or pollutants which reflects its integrity by creating lesions and other histopathological alterations of the liver parenchyma.<sup>[3]</sup> They are used as popular experimental tools for the investigation of liver injury. The liver damage produced by these agents is seen shortly after the exposure to the toxic substances and is dose related.<sup>[4]</sup> Having analysed the biochemical parameters in order to assess the response of the organs to oxidative stress, a treatment with the leaf extract, the tissues (liver slices) were also examined for histopathological changes like necrosis, edema and lymphocyte infiltration in the presence of the oxidant (H<sub>2</sub>O<sub>2</sub>) with and without the leaf extract. The procedure of Luna 1968<sup>[5]</sup> was followed for this study. The hepatic lobule is defined histologically as a hexagonal region of parenchyma which surrounds the central vein at its centre.

**MATERIALS AND METHODS**

The tissues were placed in 10% formal saline (10% formalin in 0.9% NaCl) for one hour. They were then left overnight in running water after securing the mounts of the vessels with cotton gauze. The tissues were dehydrated in ascending grade of isopropanol by immersing in 80% isopropanol overnight followed by 100% isopropanol for one hour. The dehydrated tissues were cleared in two changes of xylene, one hour each. Then the tissues were impregnated with histology grade paraffin wax at 60°C. The impregnated tissues were embedded in paraffin blocks using the same grade wax. The paraffin blocks were mounted and cut with a rotary microtome at 3 micron thickness. The sections were floated on a tissue mount and cut with a rotary microtome at 3 micron thickness. The sections were floated on a tissue floatation bath at 40° C and taken on a glass slide smeared with equal parts of egg albumin and glycerol. The sections were then melted in an incubator at 60°C and after 5 minutes allowed to cool.

**Tissue staining:** The sections were deparaffinized by immersing in xylene for 10 minutes in a staining jar. The deparaffinized sections were washed in 100% isopropanol and stained in Ehrlich's hematoxylin for 8 minutes. After staining in hematoxylin, the sections were washed in tap water and dipped in a alcohol (8.3% HCl in 70% alcohol) to remove excess stain. The section stained in 1% aqueous solution of eosin for 1 minute. The excess stain was washed in tap water and the sections were allowed to dry. The complete dehydration

of the stained sections were ensured by placing the sections in the incubator at 60°C for 4 minutes. When the sections were cooled, they were mounted in DPX mountant. The cell architecture in the liver was observed under high power objective in a microscope.


**Groups used for the study:** Four different groups were studied to determine the protective effect of the methanolic extract of the *M.hortensis* leaves. First was the control which contains only the liver extract. Second contains only the methanolic extract of the leaf (1g/mL). The third treatment group contains the oxidant namely H<sub>2</sub>O<sub>2</sub> (10%) and the fourth group is exposed to the oxidant and also the leaf extract to analyze the extent of protection rendered by the leaf in the presence of the oxidant.

## RESULTS


Histopathological analysis was carried out to confirm the spectrophotometric analysis. In the control group, the liver tissue showed normal architecture with intact portal triad. In the presence of the oxidant, the liver tissue showed cellular edema. This is presumably due to the presence of stress, which caused the normal cells to swell and exhibit edema thereby blocking the sinusoids or minimizing the sinusoidal spaces. The liver tissue in the presence of the plant extract alone showed a lower extent of cellular edema.

Untreated goat liver slices showed normal hepatic architecture and no fatty changes (Fig.1a). Administration of H<sub>2</sub>O<sub>2</sub> resulted in fatty change and


lobular ballooning followed by degeneration of hepatocytes. Administration of the methanolic leaf extract of *M. hortensis* exhibited significant improvement (Fig.1b). The hepatocytes showed only mild fatty change. The liver showed only mildly distorted architecture. Central vein showed no distinct changes. Portal triad exhibited only mild fibrosis with mild inflammation. Histological profile of the liver slices exposed to the oxidant alone reported to show intense centrilobular necrosis, steatosis, swelling of hepatic cytoplasm, ballooning degeneration, fatty changes and broad infiltration of lymphocytes and Kupffer cells around the central vein (Fig. 1c). When the oxidant was co-administered along with the leaf extract, notable damage was observed in certain areas, but some parts of the liver tissue showed recovered areas with normal architecture. The periportal areas showed some edema but the peripheral areas showed preserved architecture. This indicated that the methanolic extract of the *M. hortensis* leaves is effective in protecting the liver tissue from oxidative damage. The liver showed distorted architecture with nodule formation, distorted central vein and the portal triad showed fibrous portal expansion with moderate fibrosis and moderate inflammation (Fig. 1d). Histological observations basically supported the results obtained from the other biochemical investigations. The mechanism of action of the plant remains to be studied. The above results support the incorporation of *M. hortensis* as a component with protective effect against the action of oxidative stress by hydrogen peroxide.




**Fig. 1a: Untreated Liver Slice.**



**Fig. 1b: Leaf Extract.**



**Fig. 1c: Oxidant.**



**Fig. 1d: Oxidant + Leaf Extract.**

**Figure 1: Histopathological Architecture of the goat liver slices.**

## DISCUSSION

Extrapolations from the results obtained in precision cut liver slices to the *in vivo* condition have been successfully established for metabolic clearance, metabolism, and toxicity of several drugs.<sup>[6]</sup> The goat liver slices exposed to oxidant in the presence and absence of the leaf extract were analyzed histopathologically and the results were in agreement to those obtained with the antioxidant status of the slices. The presence of the oxidant caused tissue damage, which was reverted back to a certain extent due to the presence of the leaf extract. There is a rich source of literature that adds credibility to the fact that histopathological analysis supports all *in vitro* studies carried out in cell free system. It was suggested<sup>[7]</sup> that careful histopathologic evaluation and documentation of all manipulations in any model are needed if progress in the field is to occur and be carried into human studies. Tenpe<sup>[8]</sup> concluded that the *Oroxylum indicum* leaf extract was a good antioxidant and expressed hepatoprotective activity histologically. Ali<sup>[9]</sup> demonstrated the protective ability of bark of *Khaya senegalensis* on liver injury induced by CCl<sub>4</sub> in albino rats. It was indicated<sup>[10]</sup> the protective role of methanolic extract of *Digera muricata* on levels of antioxidants in rat thyroid against the toxicity of CCl<sub>4</sub>. The protective effects of *Hellinu linteus* on CCl<sub>4</sub> induced rat liver damage showed CCl<sub>4</sub> induced histological changes like increased degeneration, necrosis, hepatitis and portal triaditis.<sup>[11]</sup> The protective effect of the *Moringa oleifera* leaf extract against CCl<sub>4</sub> induced oxidative stress in precision cut liver slices was confirmed by histopathology analysis.<sup>[12]</sup>

## CONCLUSION

Thus it can be inferred from the results obtained from the histological architecture that the methanolic extract of *M.hortensis* protected the liver from the histological damage by H<sub>2</sub>O<sub>2</sub>. Also, the histopathology study has shown similar results as observed using other parameters. This indicates that the precision cut liver slices can be used for histopathology study as an alternative for *in vitro* studies which can yield reproducible results.

## ACKNOWLEDGEMENT

The authors are grateful to the WoS-A (Women Scientist Scheme A) scheme, Department of Science and Technology, New Delhi for their financial support. Also to Department of Histopathology, G.Kuppuswamy Naidu Memorial Hospital, Coimbatore, Tamil Nadu for interpreting the histopathology results.

## REFERENCES

1. Palaniswamy R, Padma PR. Antioxidant effect of *Majorana hortensis* leaves. International Journal of Research in Ayurveda and Pharmacy, 2011; 2(4): 1365-1367.

2. Eide M, Karlsen QA, Kryvi H, Olsvik PA, Goksoyr A. Precision cut liver slices of Atlantic cod: A *in vitro* system for studying the effects of environmental contaminants. Aquatic Toxicology, 2014; 153: 110-115.
3. Roberts RJ., The pathophysiology and a systematic pathology of teleost. In: Fish pathology, Ed. R.J. Roberts Bailieve Tindal, London, 1978; 55-91.
4. Rao JK, Madhu Ch., Rama Murthy K. Histopathological and histochemical changes under phosphomidon intoxication in liver of freshwater fish *Tilapia mossambica*. Proceedings from Bulletin of Environmental Science, 1985; 3: 20-23.
5. Luna LC. Manual of Histological Staining Methods For The Armed Forces, Institute of Pathology, 3<sup>rd</sup> Edition, New York; McGraw Hill Cook Co, 1968; 1-31.
6. Olinga P. Schuppan D. Precision cut liver slices: A tool to model the liver *ex vivo*. Journal of Hepatology, 2013; 58: 1252-1253.
7. Brunt EM. Do you see what I see? Role of Quality histopathology in scientific study. Hepatology, 2008; 47: 771-774.
8. Tenpe CR, Upaganlawar A, Burle S, Yeole P. *in vitro* antioxidant and preliminary hepatoprotective activity of *Oroxylum indicum* leaf extracts. Pharmacology Online, 2009; 1: 35-43.
9. Ali SAM., Elbadwi SMA, Idris TM, Osman KM. Hepatoprotective activity of aqueous extract of *Khaya senegalensis* bark in rats. Journal of Medicinal Plants Research, 2011; 5: 5863-5866.
10. Khan AM, Khan GN, Shabbir M, Saeed N. Shah NA, Bokhara J and Rashid U. Protective effects of *Digera muricata* (L.) mart. against carbon tetrachloride induced oxidative stress in thyroid of rat. African Journal of Biotechnology, 2011; 10: 175754-175760.
11. Huang G, Deng JS, Chiu CS, Liao JC, Hsieh W, Shewu M, Wu, C. Hispolon protects against acute liver damage in the rat by inhibiting lipid peroxidation, proinflammatory cytokine and oxidative stress and downregulating the expressions of iNos, COX-2 and MMP-p, eCAM, Article ID 480714, 2012; 12. DOI:10.1155/2012/480714
12. Sreelatha S., Padma P.R. Protective Mechanism of *Moringa olifera* against CCl<sub>4</sub> induced oxidative stress in precision cut liver slices. Forsch kplementmed, 2011; 17: 189-194.