

## ROLE OF PROBIOTICS IN MANAGEMENT OF NEONATAL JAUNDICE

Dr. Khadija Khan\*, Dr. Haseeb Iftikhar, Dr. Muhammad Awais

Pakistan.

\*Corresponding Author: Dr. Khadija Khan

Pakistan.

Article Received on 13/03/2018

Article Revised on 03/04/2018

Article Accepted on 24/04/2018

## ABSTRACT

**Objective:** To evaluate the therapeutic effects of probiotics on neonatal jaundice and the safety. **Methods:** Sixty-eight neonates with jaundice were divided into a control group and a treatment group (n=34) randomly, and treated by blue light phototherapy and that in combination with probiotics. The serum bilirubin levels were detected before and 1, 4, 7 days after treatment. The time when therapy showed effects and jaundice faded, clinical outcomes as well as adverse reactions were recorded. The categorical data were expressed as ( $\pm$  s) and compared by t test. The numerical data were expressed as (case, %) and compared by  $\chi^2$  test.  $P < 0.05$  was considered statistically significant. **Results:** Serum bilirubin levels of the two groups were similar before treatment ( $P > 0.05$ ). The levels significantly decreased 1, 4 and 7 days after treatment ( $P < 0.05$ ), but there was no significant inter-group difference on the post-treatment 1st day ( $P > 0.05$ ). The treatment group underwent more significant decreases on the 4th and 7th days than the control group did ( $P = 0.002, 0.001$ ). In the treatment group, the therapy exerted effects on ( $1.0 \pm 0.5$ ) d and jaundice faded on ( $3.8 \pm 1.7$ ) d, which were ( $2.6 \pm 0.6$ ) d and ( $5.3 \pm 2.1$ ) d respectively in the control group ( $P = 0.001, 0.002$ ). The effective rate of the treatment group significantly exceeded that of the control group ( $P = 0.002$ ). There were no obvious adverse reactions in either group. **Conclusions:** Probiotics lowered the serum bilirubin levels of neonates with jaundice rapidly, safely and significantly, and accelerated jaundice fading as well. This method is worthy of application in clinical practice.

**KEYWORDS:** Bifid triple viable, Jaundice, Neonate, Probiotics.

## INTRODUCTION

In clinical practice, neonatal jaundice is very common, which is mainly manifested as yellowish pigmentation of mucous membranes due to hyperbilirubinemia.<sup>[1]</sup> The incidence of neonatal jaundice remains high because of infections, genetic factors and premature birth. If not timely treated, neonates may be subjected to pneumonia or septicemia that severely endangers normal development of intelligence, hearing and the nervous system, or leads to permanent sequelae.<sup>[2]</sup>

Besides blue light phototherapy, touching and drugs, probiotics have also been used to enhance immunity mainly by regulating bacterial colonies. They can form a biological barrier by specifically binding intestinal epithelial cells through teichoic acid. Therefore, particular attention has been paid to the use of probiotics in treating neonatal jaundice. Therefore, the aim of this study was to evaluate the therapeutic effects of probiotics on neonatal jaundice and the safety, providing clinical evidence for future treatment.

## METHODS

**General Information:** Sixty-eight neonates with jaundice who were treated in Children hospital Lahore from July 2012 to August 2014 were selected in this study. This study was approved by the ethics committee of our hospital. Written consent has been obtained from the legal guardians of these neonates.

They were diagnosed according to the criteria for neonatal jaundice based on the transcutaneous bilirubin levels ( $> 12.9$  mg/dL, performed by NJ33A infants tester). Then they were randomly divided into a control group and a treatment group (n=34).

The control group comprised 20 boys and 14 girls with the gestational ages of 38-42 weeks (average:  $39.5 \pm 0.3$ ). They were aged 5-18.2 d, ( $11.2 \pm 1.3$ ) d in average. Their birth weights ranged from 3.12 to 4.21 kg, ( $3.75 \pm 0.21$ ) kg in average. Causes of jaundice: 10 cases of hemolytic jaundice, 8 cases of infectious jaundice, 8 cases of dysmetabolic jaundice and 8 cases of breast milk jaundice. The treatment group consisted of 22 boys and 12 girls with the gestational ages of 37-41 weeks (average:  $39.0 \pm 0.4$ ). They were aged 4.2-18.5 d, ( $11.3 \pm 1.2$ ) d in average. Their birth

weights ranged from 3.20 to 4.37 kg, (3.86±0.23) kg in average. Causes of jaundice: 8 cases of hemolytic jaundice, 11 cases of infectious jaundice, 8 cases of dysmetabolic jaundice and 7 cases of breast milk jaundice. The two groups had similar age, gender ratio and pretreatment bilirubin levels (P>0.05).

**Treatment Methods:** The control group was treated by blue light phototherapy for 3-5 d, 12-24 hour per day. Meanwhile, they were intravenously infused with phenobarbital.

Based on these, the treatment group was also orally administered with bifid triple viable

**Observation Indices:** Fasting venous bloods (3 ml) were drawn before and 1, 4 and 7 days after treatment to detect serum bilirubin levels. After centrifugation at 3000 r/minutes for 10 minutes, the serum was collected and detected by using Mindray BS400 Automatic Biochemical Analyzer with the oxidation method.<sup>[4]</sup> The time when therapy showed effects and jaundice faded, clinical outcomes and adverse reactions were recorded. Time of exerting effects: The time of icteric index that was measured with a transcutaneous jaundice meter decreased.<sup>[5]</sup>

**Clinical outcomes: Markedly effective:** Fading of yellowish pigmentation and obvious decrease in serum bilirubin level. Effective: Basic fading of

**Table-II: Changes of clinical indices (±s).**

Group	Time of exerting effect (d)	Time of jaundice fading (d)	No. of blue light phototherapy	Time of blue light phototherapy (h)
Control (n=34)	2.6±0.6	5.3±2.1	2.69±0.96	53.12±19.67
Treatment (n=34)	1.0±0.5	3.8±1.7	2.17±0.76	39.74±9.57
t	3.750	2.121	2.384	3.059
P	0.001	0.002	0.001	0.004

D. respectively in the control group (P<0.05) In addition, the two groups had significantly different numbers and times of blue light phototherapy (P<0.05) (Table-II).

**Clinical Outcomes:** The effective rate of the treatment group significantly exceeded that of the control group (P<0.05) (Table-III).

**Adverse Reactions:** Neither group was subjected to obvious adverse reactions, indicating the treatment was safe.

## DISCUSSION

Neonatal jaundice, which is common in clinical practice, can mainly be classified as physiological and pathological ones. The former does not need

yellowish pigmentation and moderate decrease in serum bilirubin level. Ineffective: Unchanged yellowish pigmentation and maintained or elevated serum bilirubin level.

## RESULTS

**Serum Bilirubin Levels:** Serum bilirubin levels of the two groups were similar before treatment (P>0.05). The levels were significantly decreased 1, 4 and 7 days after treatment (P<0.05), but there was no significant inter-group difference on the post-treatment 1st day (P>0.05). The treatment group underwent more significant decreases on the 4th and 7th days than the control group did (P<0.05) (Table-I).

**Clinical Indices:** In the treatment group, the therapy exerted effects on (1.0±0.5) d and jaundice faded on (3.8±1.7) d, which were (2.6±0.6) d and (5.3±2.1)

**Table I: Serum bilirubin levels before and after treatment (±s, μmol/L).**

Group	before	1 d after	4 d after	7 d after
Control (n=34)	351±48	312±41a	195±40a	108±21a
Treatment (n=34)	347±52	309±42a	155±37a	79±18a
T	0.367	0.876	2.747	5.918
P	>0.05	>0.05	0.002	0.001

Compared with the levels before treatment, aP<0.05.

special treatment,<sup>[6]</sup> but the latter, which originates from various factors, easily leads to bilirubin encephalopathy and even brain damage or death. Therefore, it should be timely treated. Mainly manifested as hyperbilirubinemia, neonatal jaundice is caused by enhanced intestinal-hepatic circulation of bilirubin owing to high content and activity of β-glucuronidase (β-GD).<sup>[7]</sup> β-GD can hydrolyze the bound bilirubin into unbound one and glucuronide, and the unbound bilirubin boost intestinal-hepatic circulation after being absorbed by intestinal cells.<sup>[8]</sup> Recently, intestinal flora, which are diverse and structurally complex, have attracted wide attention as the "Second Human Genome".<sup>[9]</sup>

The roles of probiotics in human body have been explained by different pharmacological mechanisms. For example, they can rapidly increase

the number of anaerobic bacterial colonies, promote the recovery of intestinal microflora balance, and resist infections in some cases.<sup>[10]</sup> On the other hand, intestinal probiotics affect the amount of bilirubin in the enterohepatic circulation by reducing the degradation of bound bilirubin.<sup>[11]</sup>

Group	Markedly Effective	Effective	Ineffective	Effective rate (%)
Control (n=34)	18	9	7	79.41
Treatment (n=34)	23	8	3	91.18
$\chi^2$				10.65
P				0.002

Meanwhile, they are able to stimulate intestinal peristalsis, which also benefits the elimination of bilirubin. It has previously been reported that oral administration of probiotics showed markedly better effects than those of routine blue light phototherapy.<sup>[12]</sup> Furthermore, the establishment of intestinal probiotic bacteria can perfect the human immune system, thereby affecting the activation and proliferation of sIg A and T lymphocytes as an analogue to active immunity.<sup>[13,14]</sup> As a result, these exert obvious beneficial effects on neonates.<sup>[15]</sup>

In this study, neonatal jaundice was treated by using bifid triple viable containing *Lactobacillus bulgaricus*, live *Bifidobacterium* and *Streptococcus thermophilus* that are normal microorganisms in the human intestinal tract.<sup>[16]</sup> After being orally taken, they grew in the intestinal tract to produce vitamins and to help the proliferation of normal bacterial colonies. Serum bilirubin levels of the two groups were similar before treatment ( $P>0.05$ ). The levels were significantly reduced 1, 4 and 7 days after treatment ( $P<0.05$ ), but there was no significant inter-group difference on the post-treatment 1st day ( $P>0.05$ ). The treatment group underwent more significant decreases on the 4th and 7th days than the control group did ( $P<0.05$ ). In the treatment group, the therapy exerted effects on  $(1.0\pm 0.5)$  d and jaundice faded on  $(3.8\pm 1.7)$  d, which were  $(2.6\pm 0.6)$  d and  $(5.3\pm 2.1)$  d respectively in the control group, with significant differences ( $P<0.05$ ). Moreover, the two groups had significantly different numbers and times of blue light phototherapy ( $P<0.05$ ).

Accordingly, bifid triple viable accelerated jaundice fading by rapidly lowering the bilirubin level, with quick recovery also. Probably, this agent in the intestinal tract of neonates, and the resulting metabolites effectively corrected the slightly alkaline environment, thus weakening the activity of  $\beta$ -GD and preventing it from binding and hydrolyzing bilirubin.<sup>[17]</sup> Furthermore, the activities of liver enzymes were also enhanced, which benefited the excretion and binding of bilirubin.<sup>[18]</sup> The effective rate of the treatment group was significantly higher

than that of the control group ( $P<0.05$ ). Since neither group was subjected to obvious adverse reactions, the treatment was fairly safe.

In summary, probiotics were able to treat neonatal jaundice quickly, safely and significantly, without discernible side effects. Hence, this method is worthy of application in clinical practice.

**Declaration of interest:** The authors have no Conflicts of interest to declare.

**Source of funding:** None.

## REFERENCES

- Porter ML, Dennis BL. Hyperbilirubinemia in the term newborn. *Am Fam Physician*, 2002; 65(4): 599-606.
- Bjarnason I, MacPherson A, Hollander D. Intestinal permeability: an overview. *Gastroenterology*, 1995; 108(5): 1566-1581.
- Hu YM, Jiang ZF Eds. *Zhu Futang Practice of Pediatrics*. 7<sup>th</sup> Edition. People's Medical Publishing House Co., Ltd., 2002.
- Wolff M, Schinasi DA, Lavelle J, Boorstein N, Zorc JJ. Management of neonates with hyperbilirubinemia: Improving timeliness of care using a clinical pathway. *Pediatrics*, 2012; 130(6): e1688-1694. doi: 10.1542/peds.2012-1156.
- Salinas M, López-Garrigós M, Lugo J, Gutiérrez M, Flors L, Leiva-Salinas C. Diagnostic accuracy of icteric index to detect abnormal total bilirubin values. *J Clin Pathol*. 2012; 65(10): 928-933. doi: 10.1136/jclinpath-2012-200811.
- Bertini G, Dani C Eds. *Bilirubin Metabolism, Unconjugated Hyperbilirubinemia, Physiological Neonatal Jaundice*. Neonatol. Springer-Verlag Milan, 2012; 608-610.
- Raju TN. Developmental physiology of late & moderate prematurity. *Semin Fetal Neonatal Med*, 2012; 17(3): 126-131. doi: 10.1016/j.siny.2012.01.010.
- Wang X, Chowdhury JR, Chowdhury NR. Bilirubin metabolism: Applied physiology. *Curr Pediatr*, 2006; 16(1): 70-74. doi: 10.1016/j.cupe.2005.10.002.
- Zheng SF, Xu YZ, Ai Q, Yu JL. Intestinal microbial community diversity in neonatal necrotizing enterocolitis analyzed by PCR-DGGE. *Chongqing Medicine*, 2012; 41(29): 3026-3029.
- Chen CH, Wu Q, Li MJ. [Management of Hyperbilirubinemia in Newborns and Related Issues]. *Journal of Applied Clinical Pediatrics*, 2011; 14(26): 1132.
- Caffarelli C, Santamaria F, Vottero A, Dascola CP, Mirra V, Sperli F, et al. Progress in pediatrics in 2013: choices in allergology, endocrinology, gastroenterology, hypertension,

- infectious diseases, neonatology, neurology, nutrition and respiratory tract illnesses. *Ital J Pediatr.* 2014; 40(1): 62. doi: 10.1186/1824-7288-40-62.
12. Jia HY. Effect of oral probiotics on treatment of neonates with hyperbilirubinemia and its influence on immune function. *J Clin Med Pract,* 2015; 19(7): 104-106.
  13. Stevenson DK, Wong RJ. Metalloporphyrins in the management of neonatal hyperbilirubinemia. *Semin Fetal Neonatal Med.,* 2009; 15(3): 164-168. doi: 10.1016/j.siny.2009.11.004.
  14. Bengmark S. Ecnutrition and health maintenance-a new concept to prevent GI inflammation, ulceration and sepsis. *Clin Nutr,* 1996; 15(1): 1-4.
  15. Rayes N, Seehofer D, Theruvath T, Mogl M, Langrehr JM, Nüssler NC, et al. Effect of enteral nutrition and synbiotics on bacterial infection rates after pylorus-preserving pancreatoduodenectomy: randomized, double-blind trial. *An Surg,* 2007; 246(1): 36-41. doi: 10.1097/01.sla.0000259442.78947.19.
  16. Li G, Zeng S, Liao W, Lv N. The effect of bifid triple viable on immune function of patients with ulcerative colitis. *Gastroenterol Res Pract,* 2012; 2012: 404752. doi:10.1155/2012/404752.
  17. Maldonado J, Cañabate F, Sempere L, Vela F, Sánchez AR, Narbona E, et al. Human milk probiotic *Lactobacillus fermentum* CECT5716 reduces the incidence of gastrointestinal and upper respiratory tract infections in infants. *J Pediatr Gastroenterol Nutr,* 2012; 54(1): 55-61. doi: 10.1097/MPG.0b013e3182333f18.
  18. Minomo A, Ishima Y, Chuang VT, Suwa Y, Kragh-Hansen U, Narisoko T, et al. Albumin domain II mutant with high bilirubin binding affinity has a great potential as serum bilirubin excretion enhancer for hyperbilirubinemia treatment. *Biochim Biophys Acta,* 2013; 1830(4): 2917-2923. doi: 10.1016/j.bbagen.2013.01.006.