

**FREQUENCY OF MICROORGANISMS CAUSING URINARY TRACT INFECTION  
AMONG DIABETIC PATIENTS**Dr. Maha Saher\*<sup>1</sup>, Dr. Aliza Abbass Syed<sup>2</sup> and Dr. Hamna Javaid<sup>3</sup><sup>1</sup>Pmdc#88804-p.<sup>2</sup>Pmdc#80318-p.<sup>3</sup>Pmdc#88363-p.

\*Corresponding Author: Dr. Maha Saher

Pmdc#88804-p.

Article Received on 20/06/2018

Article Revised on 01/07/2018

Article Accepted on 22/07/2018

**ABSTRACT**

**Aim of study:** The prevalence of asymptomatic and symptomatic UTI is very high among diabetic patients. This study was conducted to assess the frequency of microorganism involved in urinary tract infection among diabetic patients. **Methodology:** It is cross-sectional study done at Pathology department and Diabetic OPD of Jinnah Hospital, Lahore for a period of 6 months from December 2016 to April 2017. Total 250 samples of Urine was collected from patients visiting diabetic OPD. Patients having symptoms of UTI or having more than 5 WBC/HPF on urine analysis were selected for the study and urine culture was performed on their samples. Using a standard quantitative loop, urine samples (1  $\mu$ L and 10  $\mu$ L) were used to inoculate Cysteine lactose electrolyte deficient (CLED) agar (Oxoid, Basingstoke, UK), MacConkey, 5% Sheep Blood agar, and chromogenic UTI (Oxoid) agar plates. Plates were incubated for 24 h at 37°C and the outcome was judged as significant/nonsignificant growth, or contaminated (discarded). Significant bacteriuria was defined as urine culture plates showing  $\geq 10^5$  colony-forming units (CFU)/mL of single bacterial species. **Results:** Total 250 urine samples of diabetic patients were selected for study. Among these patients 162 (64.8%) were female and 88 (35.2%) were males. 171 (68.4%) patients were having symptoms of urinary tract infection and remaining 79 (31.6%) were asymptomatic. Asymptomatic patients were having more than 5 WBC/HPF on complete urine analysis. 31 patients were having HbA<sub>1c</sub> with in target range (i.e. 6-7), 100 patients were having HbA<sub>1c</sub> 7-8 and remaining 119 patients were having HbA<sub>1c</sub> above 8. After 24 hours of intubation in the required media we found 20 sample were having mixed growth/contaminated, 195 (78%) sample were having E.coli growth, in 64 (25.6%) patients were Klebsiella spp, Staphylococcus in 31 (12.4%), Pseudomonas in 15 (6%), Proteus in 9 (3.6%), Streptococcus in 2 (0.8%), Enterococcus spp. in 3 (1.2%) and Candida spp. in 1 (0.4%). **Conclusion:** UTIs are frequent in patients with diabetes. The most frequent microorganism is *E. coli* and fungal infections are least common in diabetic patients in our study. Symptomatic UTI are more common in patients due to late presentation to hospital. Among asymptomatic UTI cases, female were more common as compared to male.

**KEYWORDS:** UTI, Diabetes mellitus, CLED, E, coli, asymptomatic, symptomatic.**INTRODUCTION**

Diabetes mellitus has become a major health challenge worldwide. As the prevalence of diabetes mellitus increases worldwide, complications associated with it, also have equal importance. Diabetes mellitus has long been considered to be a predisposing factor for urinary tract infection (UTI),<sup>[1]</sup> and the urinary tract is the principle site of the infection in diabetic patients with increased risk of complications of UTI such as emphysematous cystitis, severe urosepsis, pyelonephritis and renal papillary necrosis etc.<sup>[2]</sup> The mechanisms which potentially contribute to UTI in these patients are impairments in the immune system,<sup>[3,4]</sup> poor metabolic control of diabetes,<sup>[5,6]</sup> and incomplete bladder emptying due to autonomic neuropathy.<sup>[7,8]</sup> Factors that were found

to enhance the risk for UTI in diabetics include age, metabolic control, and long term complications, primarily diabetic nephropathy and cystopathy.<sup>[9]</sup> Females are more prone to UTIs due to short urethra, pregnancy and sexual activity. In pregnancy most females develops gestational diabetes which leads to UTIs. Canadian and Danish studies shows diabetics were mostly hospitalized with urinary tract infections.<sup>[10]</sup> Asymptomatic bacteriuria (ASB) and incidence of urinary tract infections (UTIs) is high in diabetic patients.<sup>[11]</sup> The most common cause of UTI in men and women with and without DM is *Escherichia coli*. According to a study the organisms causing UTI in diabetic female are *Escherichia coli* 54.1%, *Enterococcus* spp 8.3%, and *Pseudomonas* spp 3.9%.<sup>[1]</sup>

The diagnosis of UTI should be suspected in any diabetic patient if these symptoms are present e.g. frequency, urgency, dysuria, and suprapubic pain, costovertebral angle pain/tenderness, fever, and chills. A urine culture should be obtained in all cases of suspected UTI in diabetic patients, prior to initiation of treatment. The preferred method of obtaining a urine culture is from voided, clean-catch, midstream urine.<sup>[12]</sup> Increase in incidence of urinary tract infection due to increase in diabetic patients worldwide, may impose a substantial burden on medical costs.<sup>[13]</sup> In addition, the high rates of antibiotic prescription, including broad-spectrum antibiotics, for UTI in these patients may further induce the development of antibiotic-resistant urinary pathogens.<sup>[14]</sup> The Objective of current study is to assess the type of microorganisms and their antimicrobial susceptibility in diabetic patients at a tertiary care unit.

## METHODOLOGY

It is cross-sectional study done at Pathology department and Diabetic OPD of Jinnah Hospital, Lahore for a period of 6 months from December 2016 to April 2017. Total 250 samples of Urine was collected from patients visiting diabetic OPD. Sample was collected in wide mouth leak proof container. Mid-stream urine sample (15-20ml) collected from non-catheterized patients. Labeled sample along with filled pre-formed questionnaire deposited to Pathology Lab within an hour. Informed consent was taken from the candidates.

### Inclusion criteria

Patients having symptoms of UTI or having more than 5 WBC/HPF on urine analysis were selected for the study and urine culture was performed on their samples.

### Exclusion criteria

Patients having pregnancy, any renal pathology, took antibiotics within last 15 days, history of any surgery in recent days, or having any other infection e.g.

respiratory, skin or mucous membrane were selected in the study.

Using a standard quantitative loop, urine samples (1  $\mu$ L and 10  $\mu$ L) were used to inoculate Cysteine lactose electrolyte deficient (CLED) agar (Oxoid, Basingstoke, UK), MacConkey, 5% Sheep Blood agar, and chromogenic UTI (Oxoid) agar plates. Plates were incubated for 24 h at 37°C and the outcome was judged as significant/nonsignificant growth, or contaminated (discarded). Significant bacteriuria was defined as urine culture plates showing  $\geq 10^5$  colony-forming units (CFU)/mL of single bacterial species. Statistical analysis was done by using SPSS version 20.0.

## RESULTS

Total 250 urine samples of diabetic patients were selected for study. Among these patients 162 (64.8%) were female and 88 (35.2%) were males. 171 (68.4%) patients were having symptoms of urinary tract infection and remaining 79 (31.6%) were asymptomatic. Asymptomatic patients were having more than 5 WBC/HPF on complete urine analysis. 31 patients were having HbA<sub>1c</sub> with in target range (i.e. 6-7), 100 patients were having HbA<sub>1c</sub> 7-8 and remaining 119 patients were having HbA<sub>1c</sub> above 8. After 24 hours of intubation in the required media we found 20 sample were having mixed growth/contaminated, 195 (78%) sample were having E.coli growth, in 64 (25.6%) patients were Klebsiella spp, Staphylococcus in 31 (12.4%), Pseudomonas in 15 (6%), Proteus in 9 (3.6%), Streptococcus in 2 (0.8%), Enterococcus spp. in 3 (1.2%) and Candida spp. in 1 (0.4%).

**Table 1: gender of 250 samples selected for the study.**

Gender	Number of patients	Percentage
Male	88	35.2%
Female	162	64.8%

**Table 2: number of patients presented with symptomatic or asymptomatic UTI.**

	Number of patients	Percentage
Symptomatic UTI	171	68.4%
Asymptomatic UTI	79	31.6%

**Table 3: diabetes status on the basis of HbA<sub>1c</sub> value among these 250 cases.**

Diabetes status	Number of patients	Percentage
HbA <sub>1c</sub> range 5.6-6.5 (controlled)	31	12.4%
HbA <sub>1c</sub> range 6.5-8 (uncontrolled)	100	40%
HbA <sub>1c</sub> above 8 (uncontrolled)	119	47.6%

Microorganism	Number of sample	Percentage
E. Coli spp.	195	78%
Klebsiella spp.	64	25.6%
Staphylococcus spp.	31	12.4%
Pseudomonas spp.	15	6%
Proteus spp.	9	3.6%
Streptococcus spp.	2	0.8%

Enterococcus spp.	3	1.2%
Candida spp.	1	0.4%
Contaminated/mixed growth samples	20	8%

## DISCUSSION

The ratio of symptomatic patients was higher than asymptomatic bacteriuria in this study as compared to a study reported in Ethiopia.<sup>[15]</sup> However, the frequency of ASB was higher in our study (31.6%) than previous studies (varied from 6.1% to 26.6%).<sup>[16]</sup> In our study, prevalence of UTIs in diabetic women was about 2-fold higher compared with that in diabetic men, which was related to the characteristics of female urinary tract, such as the greater length of the urethra, the greater distance between the urogenital meatus and the anus, and the antibacterial properties of the prostatic fluid.<sup>[17]</sup> Generally, compared with non-diabetic patients, diabetic patients have a higher incidence of UTI and asymptomatic bacteriuria.<sup>[18,19]</sup> The susceptibility of diabetic patients to UTI could be explained by diminished neutrophil response, lower urinary cytokines, and leukocyte concentrations, which might facilitate the adhesion of microorganisms to uroepithelial cells.<sup>[17,20]</sup> In our study we observed that with increasing the value of HbA1c, incidence of UTI increase as compared to some studies in which HbA1c seems not to be associated with the risk of UTIs.<sup>[21,22]</sup> while some studies suggested that HbA1c may be a risk factor of asymptomatic bacteriuria.<sup>[23,24]</sup> In this study, from 250 samples *E.coli* was most common (78%) followed by *Klebsiella* (25%), *Staphylococcus* spp. (12%), *Pseudomonas aeruginosa* (6%) and *Proteus* spp (3.6%). A study conducted by Bonadio et al. showed that 54.7% of UTIs were caused by *E.coli* and our findings are in accordance with these mentioned studies.<sup>[25,26]</sup> Another case-control study, conducted in New Delhi, India, that evaluated the prevalence of UTI and renal scarring in 155 patients with diabetes, also found that *E. coli* was the most commonly involved organism (64.3%), followed by *Staphylococcus aureus* (21.4%) and *Klebsiella pneumoniae* (14.3%)<sup>27</sup>. Similar findings were observed by Ramana and Chaudhary<sup>28</sup>. Bapat et al, found that *E. coli* was the most commonly isolated organism (64.3%), followed by *Staphylococcus aureus* (21.4%), and *Klebsiella pneumoniae* (14.3%).<sup>[29]</sup> Lloyds et al, have shown that *Enterococci* spp. Accounted for 35% of urinary tract isolates. We found a very low prevalence of UTI caused by fungi (*Candida* species). It is known that diabetes is a predisposing factor for fungal infections of the urinary tract. One of the most important explanations for this predisposition is glycosuria.<sup>[30]</sup> The majority of UTIs caused by fungi are clinically asymptomatic. In our study only one case showed fungal growth and that was too asymptomatic.

## CONCLUSIONS

UTIs are frequent in patients with diabetes. The most frequent microorganism is *E. coli* and fungal infections are least common in diabetic patients in our study.

Symptomatic UTI are more common in patients due to late presentation to hospital. Among asymptomatic UTI cases, female were more common as compared to male. Because of the great proportion of asymptomatic UTIs and increasing antimicrobial resistance among diabetic patients, we suggest that urine culture should be performed in all diabetic patients.

## REFERENCES

- Bonadio M, Costarelli S, Morelli G, Tartaglia T. The influence of diabetes mellitus on the spectrum of uropathogens and the antimicrobial resistance in elderly adult patients with urinary tract infection. *BMC Infect. Dis.*, 2006; 6: 54.
- Papazafiropoulou A, Daniil L, Sotiropoulos A et al. Prevalence of asymptomatic bacteriuria in Type 2 diabetic subjects with and without microalbuminuria. *BMC Research Notes.*, 2010; 3: 169-73.
- Delamaire M, Maugeudre D, Moreno M, Le Goff MC, Allannic H, Genetet B. Impaired leucocyte functions in diabetic patients. *Diabet Med.*, 1997; 14(1): 29-34.
- Valerius NH, Eff C, Hansen NE, et al. Neutrophil and lymphocyte function in patients with diabetes mellitus. *Acta Med Scand.*, 1982; 211(6): 463-467.
- Geerlings SE, Stolk RP, Camps MJ, et al. Asymptomatic bacteriuria can be considered a diabetic complication in women with diabetes mellitus. *Adv Exp Med Biol.*, 2000; 485: 309-314.
- Fünfstück R, Nicolle LE, Hanefeld M, Naber KG. Urinary tract infection in patients with diabetes mellitus. *Clin Nephrol.*, 2012; 77(1): 40-48.
- Truzzi JC, Almeida FM, Nunes EC, Sadi MV. Residual urinary volume and urinary tract infection – when are they linked? *J Urol.*, 2008; 180(1): 182-185.
- Hosking DJ, Bennett T, Hampton JR. Diabetic autonomic neuropathy. *Diabetes.*, 1978; 27(10): 1043-1055.
- Brown JS, Wessells H, Chancellor MB, et al. Urologic complications of diabetes. *Diabetes Care.*, 2005; 28(1): 177-185.
- Bonadio M, Costarelli S, Morelli G, Tartaglia T. The influence of diabetes mellitus on the spectrum of uropathogens and the antimicrobial resistance in elderly adult patients with urinary tract infection. *BMC Infect. Dis.*, 2006; 6: 54.9.
- Geerlings SE. Urinary tract infections in patients with diabetes mellitus: epidemiology, pathogenesis and treatment. *Int J Antimicrob Agents*, 2008; 31: S54-7.
- Hasan MK, Nazimuddin K, Ahmed AKMS, Sarker RSC, Haque M, Musa AKM. Differences in a bacteriological and antibiotic sensitivity patterns in

- UTI among hospitalized diabetic and non-diabetic patients. *J Medicine*, 2007; 8: 10-3.
13. Datta P, Rani H, Chauhan R, Gombar S, Chander J. Health-care-associated infections: risk factors and epidemiology from an intensive care unit in Northern India. *Indian J Anaesth*, 2014; 58(1): 30–35.
  14. Yu S, Fu AZ, Qiu Y, et al. Disease burden of urinary tract infections among type 2 diabetes mellitus patients in the US. *J Diabetes Complications*, 2014; 28(5): 621–626.
  15. Venmans LM, Hak E, Gorter KJ, Rutten GE. Incidence and antibiotic prescription rates for common infections in patients with diabetes in primary care over the years 1995 to 2003. *Int J Infect Dis.*, 2009; 13(6): e344–e351.
  16. Nicolle LE. Urinary tract infection in diabetes. *Curr Opin Infect Dis.*, 2005; 18: 49–53.
  17. Turan H, Serefhanoglu K, Torun AN, et al. Frequency, risk factors, and responsible pathogenic microorganisms of asymptomatic bacteriuria in patients with type 2 diabetes mellitus. *Jpn J Infect Dis.*, 2008; 61(3): 236–238.
  18. Boroumand MA, Sam L, Abbasi SH, Salarifar M, Kassaian E, Forghani S. Asymptomatic bacteriuria in type 2 Iranian diabetic women: a cross sectional study. *BMC Womens Health.*, 2006; 6: 4.
  19. Lipsky BA. Urinary tract infections in men. Epidemiology, pathophysiology, diagnosis, and treatment. *Ann Intern Med*, 1989; 110: 138-150.
  20. Guillausseau PJ, Farah R, Laloi-Michelin M, Tielmans A, Rymer R, Warnet A. Urinary tract infections and diabetes mellitus. *Rev Prat*, 2003; 53(16): 1790–6.
  21. Geerlings SE. Urinary tract infections in patients with diabetes mellitus: epidemiology, pathogenesis and treatment. *Int J Antimicrob Agents*, 2008; 31(1): S54–7.
  22. Soo Park B, Lee SJ, Wha Kim Y, Sik Huh J, Il Kim J, Chang SG. Outcome of nephrectomy and kidney-preserving procedures for the treatment of emphysematous pyelonephritis. *Scand J Urol Nephrol*, 2006; 40(4): 332–338.
  23. Boyko EJ, Fihn SD, Scholes D, Abraham L, Monsey B. Risk of urinary tract infection and asymptomatic bacteriuria among diabetic and nondiabetic postmenopausal women. *Am J Epidemiol*, 2005; 161(6): 557–564.
  24. Keleştimur F, Unal A, Paşaoğlu H, Başar E, Kiliç H, Doğanay M. [Asymptomatic bacteriuria in patients with diabetes mellitus]. *Mikrobiyol Bul*, 1990; 24(2): 126–132.
  25. Isenberg, H. D. Clinical microbiology procedures handbook. Washington, D.C: *American Society for Microbiology*, 1992.
  26. Collee JG, Miles RS, Watt B. Tests for identification of bacteria. In: Collee JG, Fraser AG, Marmion BP, Simmons A, editors. *Mackie and McCartney Practical Medical Microbiology*. 14th ed, New York. *Churchill Livingstone*, 2006.
  27. Bonadio M, Meini M, Spitaleri P, Gigli C. Current microbiological and clinical aspects of urinary tract infections. *Eur Urol.*, 2001; 40(4): 439-44.
  28. Bashir MF, Qazi JI, Ahmed N, Riaz S. Diversity of urinary tract pathogens and drug resistant isolates of *Escherichia coli* in different age and gender groups of Pakistanis. *Trop. J Pharm Res.*, 2008; 7: 1025-31.
  29. Ramana BV, Chaudhury A. Prevalence of uropathogens in diabetic patients and their resistance pattern at tertiary care centre in south India. *Int J Biol Med Res.*, 2012; 3(1): 1433-5.
  30. Wheat LJ. Infection and diabetes mellitus. *Diabetes Care*, 1980; 3: 187-197.