

**FREQUENCY OF MYOPIA AND UNDERLYING FACTORS OF MYOPIA AMONG  
MEDICAL STUDENT AT NISHTAR MEDICAL UNIVERSITY****Dr. Fahad Ali Khan<sup>1</sup>, Dr. Wamiq Mehmood\*<sup>2</sup>, Dr. Amr Abdullah Ahmed Shaddad<sup>3</sup>**<sup>1</sup>(PMDC # 99360-P) Medical Officer.<sup>2</sup>(PMDC # 91881-P) Medical Officer.<sup>3</sup>(PMDC # 4150-P) Medical Officer.**\*Corresponding Author: Dr. Wamiq Mehmood**

(PMDC # 91881-P) Medical Officer.

Article Received on 08/07/2018

Article Revised on 29/07/2018

Article Accepted on 19/08/2018

**ABSTRACT**

**Background:** Growing levels of high Myopia are increasing the risks of serious eye conditions which may lead to permanent blindness. A number of evaluations (at small and large scale) regarding Myopia have been carried out but very few of them reflect the opinions about medical students. **Objectives of the study:** To determine the frequency of myopia and underlying factors among 4<sup>th</sup> year students of MBBS at NMU, Multan. **Material and Methodology:** A total of 100 students at Nishtar medical university were included in study with their informed consent. A questionnaire was designed data was collected and analyzed using SPSS and Microsoft excel. Type of study was cross sectional and type of sampling was Non Probability Convenient. **Result:** The mean age of 4<sup>th</sup> year students was 22 years. Out of total 100 students, 68% were myopic. 5.9% were in the age group of 18-20 years, 83.8% in 21-23 years and 10.3% were 24-26 years of age. 66.2 % students were females. Family history of myopia was positive in 73.5% students. Genetic history of myopia is positive in 66.2%. The study hours of 19.2% students were  $\leq 2$  hour, 2-8 hours in 66.2% and  $> 8$  hours per day in 11%. The duration of TV watching was  $<2$  hour per day in 27.9% students, 2-4 hours in 41.2% and  $> 4$  hours in 30.9% children. 16.2 % of the students were spending their time on smart phones for  $<2$  hour, 58.5% for 2-4 hours and 25% for  $> 4$  hours. 8.8% students have less than 6 hours sleeping time, 64.7% students who sleep 6 to 8 hours and 26.5% more than 8 hours. **Conclusion:** From our study we concluded that Myopia is affecting majority of 4<sup>th</sup> year MBBS students. There is strong association of myopia with near work and with family and genetic history of myopia.

**KEYWORDS:** Key words are 4<sup>th</sup> year MBBS students Myopia, study hours, family history.**INTRODUCTION**

Refractive error may be defined as a state in which the optical system of the non-accommodating eye fails to bring parallel rays of light to focus on the retina. Especially myopia has become a very common problem. Myopia is a refractive error in which eye fails to see distant objects properly. It has become an ocular disorder of major public health and socioeconomic significance throughout the world. Several studies describe an increasing prevalence of myopia in the recent years. There has long been a concern that blindness and visual impairment from myopia will lead to major public health problems for many countries in Asia. Although blindness registry data indicate that myopia is the fourth leading cause of blindness in Myopia (short- or near-sightedness) affects many school-aged children and is fast becoming a major public health issue of our time. It is estimated that the current number of 2 billion people (2010) with the condition will grow to a staggering 2.6 billion by 2020 and 4.8 billion by 2050.

The onset of myopia at an early age brings with it the likelihood of life-long eye care, there is a significantly increased risk of serious ocular health problems with high myopia, which can lead to vision loss and blindness. It is projected that almost 1 billion people will have high myopia by 2050.<sup>[1]</sup>

Medical education imposes significant stress on medical students mainly through time pressure large amount of new information, excessive study hours, less sleeping hours and other problems. A considerable degree of Myopia has been reported in medical students ranging from excessive study hours to less sleeping hours and other extended near vision tasks coupled with a genetic predisposition for Myopia. This is a need to explore Myopia among medical students. Several studies suggest a high prevalence of Myopia among medical students with level of near sightedness consistently higher than in the general population and age matched peers. In fact, a recent study by National Eye Institute (NEI) shows the prevalence of Myopia grew from 25 percent of the US

Population (ages 12-54) in 1971-1972 to a whopping 41.6 percent in 1999-2004. Similarly in Norway Medical Students (University of Trondheim), of 133 (75 females, 58 males), 50.3% were found to be myopic.

## OBJECTIVES

To determine the frequency of Myopia among 4th year medical students in NMU, Multan.

To determine the underlying factor of Myopia among the students.

To assess the prevalence of Myopia between the two genders.

To assess the prevalence of Myopia among the students of different age groups.

To assess the prevalence of Myopia among the students having different genetic and family histories.

To assess the prevalence of Myopia among the students having different study and sleeping hours.

## MATERIAL AND METHODOLOGY

**Study design:** A Descriptive cross-sectional study.  
**Study Population:** 4<sup>th</sup> year MBBS at Nishtar Medical University, Multan. **Setting:** Study was carried at community medicine department at Nishtar medical University. **Study Duration:** Data was collected from 27<sup>th</sup> April 2017 to 8<sup>th</sup> May 2017. **Sampling technique:** Data was collected using Non Probability Convenient sampling. **Sample size:** 100 students from 4<sup>th</sup> year MBBS were included in the study. **Inclusion criteria:** 4<sup>th</sup> year MBBS students. **Exclusion criteria:** Students with astigmatism, hypermetropia and students not from 4<sup>th</sup> year MBBS. **Data collection procedure:** Data was collected by distributing a questionnaire that included various demographic variables like Age, Gender of the students and Residence. And various other variables suspected to be the underlying factors in causing myopia for example Study hours, sleeping hours, eye exercises, watching TV, using smart phones etc. **Data Analysis:** Data was analyzed in the department of Community Medicine using IBM SPSS (statistical package for social services) v.20. And compiling was done using MS OFFICE 2013.

**Table 7: History of myopia among the students according to genetic history.**

History of Myopia	Genetic History					
	1 Parent		2 Parent		None	
	Frequency	% age	Frequency	% age	Frequency	% age
Yes	22	32.4%	23	33.8	23	33.8
NO	9	28.1%	4	12.5	19	59.4
Total	31	31%	27	27%	42	42%

**Table 1: Age distribution of students of 4<sup>th</sup> year MBBS. n=100.**

Age group	Frequency	Percentage
18-20	7	7
21-23	81	81
24-26	12	12

**Table 2: Gender distribution of 4<sup>th</sup> year MBBS. n=100.**

Gender	Frequency	Percentage
Male	37	37
Female	63	63

**Table 3: Frequency distribution of Myopia Among 4<sup>th</sup> year MBBS students n=100.**

History of Myopia	Frequency	Percentage
Yes	68	68
No	32	32

**Table 4 Frequency distribution of Myopia among to different age groups. N=100.**

History of Myopia	Age			Total
	18-20	21-23	24-26	
Yes	4	57	7	68
No	3	24	5	32
Total	7	81	12	100

**Table 5: History of myopia according to gender distribution.**

History of Myopia	Gender		Total
	Male	Female	
Yes	23	45	68
No	14	18	32
Total	37	63	100

**Table 6: Frequency distribution of Myopia among students according to their Residence n=100.**

History of myopia	Residency		Total
	Day scholar	Boarder	
Yes	33	35	68
No	14	18	32
Total	47	53	100

Table 8: History of Myopia among the students according to Family history.

History of Myopia	Family History							
	Sibling		Grand parents		None		Both	
	Frequency	% age	Frequency	% age	Frequency	% age	Frequency	% age
Yes	34	50%	10	14.7%	18	26.5%	6	8.8%
NO	7	21.9%	5	15.6%	18	56.2%	2	6.2%
Total	41	41%	15	15%	36	36%	8	8%

Table 9: History of myopia among the students according to study hours. n=100.

History of Myopia	Study hours					
	Less than 2 hours		2 to 8 hours		More than 8 hours	
	Frequency	% age	Frequency	% age	Frequency	% age
Yes	13	19.1%	45	66.2%	10	11%
NO	10	31.2%	21	65.6%	1	3.1%
Total	23	23%	66	66%	11	14.7%

Table 10: History of myopia among the students according to Time spent in front of Display screen (laptop/pc/TV). n=100.

History of Myopia	Time Spent					
	Less than 2 hours		2 to 4 hours		More than 4 hours	
	Frequency	% age	Frequency	% age	Frequency	% age
Yes	19	27.9%	28	41.2%	21	30.9%
NO	11	34.4%	12	37.5%	9	28.1%
Total	30	30%	40	40%	30	30%

Table 11: History of myopia among the students according to Time spent on smart phone. n=100.

History of Myopia	Time Spent (TV)					
	Less than 2 hours		2 to 4 hours		More than 4 hours	
	Frequency	% age	Frequency	% age	Frequency	% age
Yes	11	16.2%	40	58.8%	17	25%
NO	5	15.6%	18	56.2%	9	28.1%
Total	16	16%	58	58%	26	26%

Table 12: History of myopia among the students according to Sleeping hours. n=100.

History of Myopia	Time Spent in sleeping					
	Less than 6 hours		6 to 8 hours		More than 8 hours	
	Frequency	% age	Frequency	% age	Frequency	% age
Yes	6	8.8%	44	64.7%	18	26.5%
NO	7	21.9%	21	65.6%	4	12.5%
Total	13	13%	65	65%	22	22%

Table 13: History of myopia among the students according to Eye exercise.

History of Myopia	Eye exercise					
	Regular		Irregular		No Exercise	
	Frequency	% age	Frequency	% age	Frequency	% age
Yes	8	11.8%	4	5.9%	56	82.4%
NO	0	0%	0	0%	32	100%
Total	8	8%	4	4%	88	88%

**Table 14: History of myopia among the students according to use of eye drops. n=100.**

History of Myopia	Use of Eye drops			
	Regular		Irregular	
	Frequency	% age	Frequency	% age
Yes	14	20.6%	54	79.4%
NO	2	6.2%	30	93.8%
Total	16	16%	84	84%

**Table 15: History of myopia among the students according to rubbing of eyes. n=100.**

History of Myopia	Rubbing of eyes			
	Yes		No	
	Frequency	% age	Frequency	% age
Yes	41	60.3%	27	39.7%
NO	21	65.6%	11	34.4%
Total	62	62%	38	38%

## DISCUSSION

In our study, out of 100 students myopia was 68% and 32% were emmetropes. Majority of students were in the age group of 21-23 years i.e. 83.8% and only 5.9% were 18-20 years, 10.3% were in 24 to 26 years. Frequency of myopia in our study was similar to myopia in the study conducted by Chaudhary *et al.*, in which it was 57.6%.<sup>[4]</sup> And 63% of the students in our RESULTS study were females while only 37% were male. There were a total of 100 students in the study. And frequency of myopia among females was higher. Frequency of myopia was 66.2% and 33.8% as compared to male, which is consistent to the study conducted by Mavracanas *TA et al.*, which reported 67.37% students were female as compared to 32.63% males.<sup>[5]</sup> Our study results revealed significant association of Family history of myopia was positive in 73.5% myopia with family history and genetic history in 66.2%. Similarly in study conducted by yingyong P parents with myopia tends to have children with myopia.<sup>[6]</sup>

According to the report by World Health Organization, uncorrected refractive error is the second commonest cause of global visual impairment next only to cataract.<sup>[1]</sup> In our study, high prevalence of myopia was found among the medical student which was in agreement with the study conducted by Sood *et al.*<sup>[2]</sup> Similar to our findings, Chalasani *et al.* also observed that the number of myopic was found to be increased among the student taking admission in the medical college every year.<sup>[3]</sup> It was observed in our study that myopia is also highly associated with use of electronic gadgets such as smart phones and computers.

This observation was in agreement with the findings of Reddy *et al.* who found that more than 2 hours continuous use of computer was significantly associated with occurrence of symptoms of computer vision syndrome.<sup>[7]</sup> Our results in this regard is also consistent with another study suggesting that prolonged use of computers is responsible for visual fatigue which in turn may lead to myopia.<sup>[8]</sup> According to David Allamby,

Founder of Focus Clinics, there has been a 35 per cent increase in the number of people with advancing myopia (short sightedness) since the launch of smart phones in 1997.<sup>[9]</sup> Our study proposed the higher incidence of myopia among smart phone users as also suggested by Lee H *et al.*<sup>[10]</sup>

## CONCLUSION

High prevalence of myopia was found among the medical students. The increased applicability of electronic gadgets, laptops, computers and smart phones were found to be the major associated risk factors along with the family history and genetic history. Further studies are recommended for the prevention of increasing frequency of myopia among the young population.

## LIMITATIONS

A very limited research has been done regarding prevalence of myopia among the 4<sup>th</sup> year MBBS students of Nishtar Medical University.

Our sample size was very small consisting of 100 students.

We used non-probability convenience sampling to draw our sample; this method is inferior to probability sampling in representation of the population and this limits the validity of the study.

Degree of generalizability is questionable.

Difficulties in estimating sampling variability and identifying possible bias.

Prospective longitudinal studies are needed to establish causal relationship between near work and myopia in the medical students.

## RECOMMENDATIONS

First of all need to identify the population at risk and myopic by screening and proper evaluation of those with

any difficulty seeing distant objects clearly such as the TV or the writing on the board. Prescribing correct (concave) glasses to individual suffering from myopia and health education of general population and high risk population regarding the underlying factors causing myopia and prevention.

Whenever close work is done

Hold the work as far away as possible.

Use as much light as possible in order to reduce the size of the pupil and, consequently, the accommodation.

Look into the distance frequently to relax the accommodation.

physical illness of smartphone users. Consumer Communications and Networking Conference (CCNC), 2013; IEEE: 821-825. DOI: 10.1109/CCNC.2013.6488555.

## REFERENCES

1. Gopalakrishnan S, Prakash MVS, Kumar R. A Study of Refractive Errors among Medical students in AIMST University, Malaysia. *Indian MedJ*, 2011; 105(11): 365.
2. Sood RS, Sood A. Prevalence of myopia among the medical students in western India visà-vis the East Asian epidemic. *IOSR JDent Med Sci.*, 2014; 13(1): 65-67.
3. Chalasani S, Jampala VK, Nayak P. Myopia among Medical Students-A Cross Sectional Study in A South Indian Medical College. *Al Ameen J Med Sci.*, 2012; 5(3): 233-242.
4. Chaudhary R, Ali H, Sheikh HN. Frequency and Ophthalmology outpatient department, it may not underlying factors of myopia among medical students be possible to generalize the results to the whole Biomedica, 2011; 27(2): 154-60.
5. Mavracanas TA, Mandalos A, Peios D, Golias V, needed to establish causal relationship between Megalou K, Gregoriadou A, et al. Prevalence of myopia in a sample of Greek students. *Acta Ophthalmol Scand* near work and myopia in this age group, 2009; 78(6): 656-59.
6. Yingyong P. Risk factors for refractive errors in primary school children (6-12 years old) in Nakhon Pathom Province. *J Med assoc*, 2010; 93(11): 1288-93.
7. Reddy SC, Low CK, Lim YP, Low LL, Mardina F, Nursaleha MP. Computer vision syndrome: a study of knowledge and practices in university students. *Nepal J Ophthalmol*, 2013; 5(10): 161-168. DOI: 10.3126/nepjoph. v5i2.8707
8. Rajeev A, Gupta A, Sharma M. Visual Fatigue and Computer Use Among College Students. *Indian JCommunity Med*, 2006; 31(3): 192-193.
9. Nnes E. Have you got 'screen sightedness'?- Smartphones are causing sight problems to soar, warns eye surgeon, "DailyMail", United Kingdom, 2013 August 15. (<http://www.dailymail.co.uk/health/article-2394611>).
10. Hosub L, Sunjae L, Young Sang C, Seo Y, Shim E. A new posture monitoring system for preventing